

A COMBINATORIAL STUDY OF AFFINE SCHUBERT VARIETIES IN THE AFFINE GRASSMANNIAN

MARC BESSON AND JIUZU HONG

ABSTRACT. Let \overline{X}_λ be the closure of the I-orbit X_λ in the affine Grassmannian Gr of a simple algebraic group G of adjoint type, where I is the Iwahori subgroup and λ is a coweight of G . We find a simple algorithm which describes the set $\Psi(\lambda)$ of all I-orbits in \overline{X}_λ in terms of coweights. We introduce R -operators (associated to positive roots) on the coweight lattice of G , which exactly describe the closure relation of I-orbits. These operators satisfy Braid relations generically on the coweight lattice. We also establish a duality between the set $\Psi(\lambda)$ and the weight system of the level one affine Demazure module $\hat{\mathcal{D}}_\lambda$ of ${}^L\tilde{\mathfrak{g}}$ indexed by λ , where ${}^L\tilde{\mathfrak{g}}$ is the affine Kac-Moody algebra dual to the affine Kac-Moody Lie algebra $\tilde{\mathfrak{g}}$ associated to the Lie algebra \mathfrak{g} of G .

1. INTRODUCTION

It is well-known that Schubert cells in the flag variety of a reductive group G can be parametrized by the elements of the Weyl group of G . Moreover, the closure relations among Schubert cells can be described by the Bruhat order on the Weyl group. There is another equivalent description of the Bruhat order in terms of the containment relations of Demazure modules of a Borel subgroup of G , which is established in the celebrated work (BGG) by Bernstein-Gelfand-Gelfand. There is a generalization of this perspective for general Kac-Moody groups, see (Ku).

From now on throughout this paper, we assume that G is a simple algebraic group over \mathbb{C} of adjoint type. Let Gr denote the affine Grassmannian $G(\mathcal{H})/G(\mathcal{O})$ of G , where \mathcal{H} is the field of Laurent series over \mathbb{C} and \mathcal{O} is the ring of formal power series. The $G(\mathcal{O})$ -orbits are indexed by dominant coweights of G . It is well-known that (cf. (BD, §4.5), (Zhu, §2.1))

$$(1) \quad \text{Gr}_\mu \subset \overline{\text{Gr}_\lambda} \text{ if and only if } \lambda - \mu \text{ is a sum of positive coroots of } G,$$

where Gr_μ and Gr_λ denote $G(\mathcal{O})$ -orbits indexed by dominant coweights λ, μ , and $\overline{\text{Gr}_\lambda}$ is the closure of Gr_λ . Moreover, the intersection cohomology of $\overline{\text{Gr}_\lambda}$ carries an action of the Langlands dual group ${}^L G$ of G , which is irreducible and of highest weight λ (cf. (MV)).

In this paper, we consider the action of the Iwahori subgroup I on Gr . The I-orbits in Gr can be indexed by coweights of G . For each coweight λ , we denote by X_λ the associated I-orbit and \overline{X}_λ the closure of X_λ in Gr . For any two coweights λ, μ , we introduce

1991 *Mathematics Subject Classification.* 05E10, 17B67, 20G44, 14M15.

Key words and phrases. Affine Schubert variety, affine Grassmannian, affine Weyl group, affine Kac-Moody algebra, level one affine Demazure module, R-operators.

the partial order $\mu <_I \lambda$ if $X_\mu \subset \bar{X}_\lambda$. This partial order is equivalent to the combinatorial Bruhat order on the affine Weyl group, see Lemma 2.22 for the precise statement. This order naturally appears in the study of parabolic Kazhdan-Lusztig polynomials for affine Kac-Moody groups (cf. (KT)), as well as in the context of non-symmetric Macdonald polynomials (cf. (Io1, Io3, Io4)). For a given coweight λ , we denote by $\Psi(\lambda)$ the set of all coweights μ such that $\mu <_I \lambda$. In this paper, we describe the partial order $<_I$ in a way similar to the condition in (1), and describe the set $\Psi(\lambda)$ by a simple algorithm. We also introduce R -operators which help to describe the extremal elements in $\Psi(\lambda)$, and they themselves satisfy braid relations generically on the weight lattice. Moreover, we give a representation theoretic interpretation of $\Psi(\lambda)$ in terms of level one affine Demazure modules, which are of twisted type when G is not simply-laced.

In Lemma 2.2 we describe an algorithm that is used to produce new elements for the set $\Psi(\lambda)$, and in Theorem 2.6 we prove that this algorithm indeed produces all elements of $\Psi(\lambda)$. Roughly speaking, any element in $\Psi(\lambda)$ can be obtained by successively adding or subtracting positive coroots, depending on the signs of the pairing between coweights and positive roots.

The set $\pi_0(\text{Gr})$ of components of Gr can be identified with the quotient group \check{P}/\check{Q} , where \check{Q} is the coroot lattice of G . Let Gr^κ be a component of Gr that contains the T -fixed point $L_{-\check{\omega}_\kappa}$ associated to the coweight $-\check{\omega}_\kappa$, where $\check{\omega}_\kappa$ is a miniscule coweight or zero. In Proposition 2.21, we explicitly realize the component Gr^κ as a partial flag variety $\tilde{G}_{\text{sc}}/\tilde{P}_\kappa$ of the affine Kac-Moody group \tilde{G}_{sc} associated to the simply-connected cover of G , and realize each I -orbit X_λ as an affine Schubert cell in $\tilde{G}_{\text{sc}}/\tilde{P}_\kappa$. In this way, we translate the partial order $<_I$ on coweights into the partial Bruhat order on W_{aff}/W_κ , where W_κ is the Weyl group of \tilde{P}_κ . In Section 2.3, we explicitly realize the affine Weyl group W_{aff} as the Weyl group of the affine Kac-Moody algebra $\tilde{\mathfrak{g}}$ associated to \mathfrak{g} , and translate the Bruhat order on W_{aff} into certain conditions on coweights (cf. Proposition 2.15, Corollary 2.12). Combining all these preparations, in Section 2.5 we prove Theorem 2.6 that is described in the previous paragraph. In Section 2.6, we use length zero elements of the extended affine Weyl group to establish bijections between the sets $\Psi(\lambda)$ in different components of the affine Grassmannian (cf. Corollary 2.25).

Let $\tilde{\mathfrak{g}}$ be the affine Kac-Moody Lie algebra associated to the Lie algebra \mathfrak{g} of G . Let ${}^L\tilde{\mathfrak{g}}$ be the affine Kac-Moody algebra with Dynkin diagram dual to that of $\tilde{\mathfrak{g}}$. As already mentioned above, the components of Gr correspond to the miniscule coweights of \mathfrak{g} or zero. Furthermore, the level one basic representations of ${}^L\tilde{\mathfrak{g}}$ also correspond to the miniscule coweights of \mathfrak{g} or zero (cf. Lemma 3.2). In summary, we have the following correspondences:

$$(2) \quad \text{Gr}^\kappa \longleftrightarrow \check{\omega}_\kappa \longleftrightarrow \mathcal{H}_\kappa,$$

where \mathcal{H}_κ denotes the associated level one basic representation of ${}^L\tilde{\mathfrak{g}}$. This suggests a duality between the affine Grassmannian Gr of G and the level one representations of ${}^L\tilde{\mathfrak{g}}$. For every coweight $\lambda \in \check{P}$, if $X_\lambda \subset \text{Gr}^\kappa$, we may associate a level one affine Demazure module $\hat{\mathcal{D}}_\lambda$ generated by a maximal weight vector $v_{\varpi(\lambda)} \in \mathcal{H}_\kappa$ associated to λ . By Proposition 3.7 together with Proposition 2.21, we can establish the following

correspondences:

$$(3) \quad \boxed{\mu <_I \lambda} \longleftrightarrow \boxed{\text{partial Bruhat order on } W_{\text{aff}}/W_\kappa} \longleftrightarrow \boxed{\hat{\mathcal{G}}_\mu \subset \hat{\mathcal{G}}_\lambda}$$

where in the first correspondence we view W_{aff} as the Weyl group of $\tilde{\mathfrak{g}}$, and in the second correspondence we view W_{aff} as the Weyl group of ${}^L\tilde{\mathfrak{g}}$ (cf. Section 3.1). Under these correspondences, in Theorem 3.9 we show that there exists a natural projection from the weight system of $\hat{\mathcal{G}}_\lambda$ to the set $\Psi(\lambda)$. It is interesting to point out that we crucially use the Frenkel-Kac construction of level one basic representations of ${}^L\tilde{\mathfrak{g}}$ in the proof of Theorem 3.9. Furthermore, in Lemma 4.7 we give a representation-theoretic interpretation for the algorithm in Lemma 2.2. More precisely we show that the algorithm of adding or subtracting positive coroots to coweights in $\Psi(\lambda)$ can be interpreted as the actions of the positive real root vectors of ${}^L\tilde{\mathfrak{g}}$ on weight vectors in $\hat{\mathcal{G}}_\lambda$.

In Section 4, we study the set $\Psi(\lambda)$ and the partial order $<_I$ further. We first show that for any two coweights $\lambda, \mu \in \check{P}$, when they are located in the same chamber, then $\mu <_I \lambda$ if and only if $\lambda - \mu$ is a sum of positive coroots relative to that chamber (cf. Theorem 4.1). This is analogous to the statement in (1). In Section 4.2, we introduce R -operators on the coweight lattice \check{P} associated to positive roots. It turns out that these operators exactly characterize the partial order $<_I$ (cf. Theorem 4.3). In Proposition 4.9 and Proposition 4.10, we describe explicitly the covering relations of $<_I$ for a coweight λ when λ is mildly regular. In Section 4.3, we show that for any two positive roots α, β , when they generate a rank two root system as simple roots, then the associated R -operators R_α, R_β satisfy a braid relation when the coweights are away from certain critical hyperplanes (Proposition 4.14). We introduce in Definition 4.5 the notion of α -regularity of a coweight in a fixed chamber, where α is a positive root. This notion allows us to cross the wall H_α defined by α so that $R_\alpha(\lambda)$ is in the reflected chamber and R_α preserves the partial order $<_I$ (cf. Proposition 3.3). This allows us to produce an algorithm to describe the vertices of the convex hull of $\Psi(\lambda)$, or in other words the moment polytope of the affine Schubert variety \bar{X}_λ , see the discussions in Section 4.4.

Acknowledgments: We would like to thank I. Cherednik, P. Fiebig and S. Kumar for helpful suggestions and discussions. We would like to thank S. Nie for bringing the reference (St) to our attentions. We also would like to thank Changlong Zhong for careful reading and pointing out some typos. J. Hong is partially supported by the Simons Foundation Collaboration Grant 524406.

2. A COMBINATORIAL DESCRIPTION OF THE CLOSURE RELATION OF IWAHORI ORBITS IN AFFINE GRASSMANNIAN

2.1. Notations. Let G be a simple algebraic group G over \mathbb{C} of adjoint type. Pick a maximal torus T contained in a Borel subgroup B of G . Let $X^*(T)$ denote the lattice of characters of T and $X_*(T)$ the lattice of cocharacters of T . Let Φ denote the set of roots for T , $\check{\Phi}$ the coroots so $(X_*(T), \Phi, X^*(T), \check{\Phi})$ is the root datum of G . Let W denote the Weyl group of G . We denote by Φ^+ (resp. $\check{\Phi}^+$) the set of positive roots (resp. positive coroots) determined by $T \subset B$. Let $\alpha_1, \dots, \alpha_\ell$ (resp. $\check{\alpha}_1, \dots, \check{\alpha}_\ell$) be simple roots

(resp. simple coroots). Under the assumption that G is of adjoint type, the lattice $X_*(T)$ coincides with the coweight lattice \check{P} .

Let $\mathcal{O} = \mathbb{C}[[t]]$ be the formal power series in t with coefficients in \mathbb{C} , and let $\mathcal{K} = \mathbb{C}((t))$ be the field of formal Laurent series in t . Let Gr denote the affine Grassmanian $G(\mathcal{K})/G(\mathcal{O})$ of G . We have an evaluation map

$$\text{ev}_0 : G(\mathcal{O}) \rightarrow G$$

sending $t \mapsto 0$. Write $\text{I} = \text{ev}_0^{-1}(B)$ as the Iwahori subgroup of $G(\mathcal{K})$. Any cocharacter $\lambda : \mathbb{C}^* \rightarrow T$ gives rise to an element $t^\lambda \in G(\mathcal{K})$. Set $L_\lambda := t^\lambda G(\mathcal{O})/G(\mathcal{O}) \in \text{Gr}$. Then all T -fixed points in Gr are given by L_λ , where $\lambda \in X_*(T)$. We denote by \mathbf{X}_λ the I -orbit $\text{I} \cdot L_\lambda$ in Gr , and we denote by Gr_λ the $G(\mathcal{O})$ -orbit $G(\mathcal{O}) \cdot L_\lambda$. The variety \mathbf{X}_λ has a unique T -fixed point L_λ , and Gr_λ has T -fixed points $L_{w(\lambda)}$, for $w \in W$.

Definition 2.1. Let $\Psi(\lambda) = \{L_\mu \in \check{P} \mid t^\mu \in \overline{\mathbf{X}_\lambda}\}$, where the closure is being taken in the Zariski topology. If $\mu \in \Psi(\lambda)$, then we write $\mu <_I \lambda$. Clearly $<_I$ gives a partial order on \check{P} .

2.2. The algorithm. We begin with a key lemma in order to describe the set $\Psi(\lambda)$.

Lemma 2.2. *Let $\lambda \in \check{P}$, and let $\alpha \in \Phi^+$ be a positive root.*

- 1) *If $\langle \lambda, \alpha \rangle > 0$, then $\lambda - k\check{\alpha} \in \Psi(\lambda)$, for $1 \leq k \leq \langle \lambda, \alpha \rangle$.*
- 2) *If $\langle \lambda, \alpha \rangle < 0$, then $\lambda + k\check{\alpha} \in \Psi(\lambda)$, for $1 \leq k \leq -\langle \lambda, \alpha \rangle - 1$.*

Proof. For any positive root $\alpha \in \Phi^+$, we may choose root subgroup homomorphisms $x_\alpha, x_{-\alpha}$ corresponding to the roots $\alpha, -\alpha$, and a cocharacter $h_\alpha : \mathbb{C}^\times \rightarrow G$, which give rise to a group homomorphism

$$\phi : \text{SL}_2 \rightarrow G$$

such that

$$\phi\left(\begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix}\right) = x_\alpha(a), \quad \phi\left(\begin{bmatrix} 1 & 0 \\ a & 1 \end{bmatrix}\right) = x_{-\alpha}(a), \quad \text{and} \quad \phi\left(\begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix}\right) = h_\alpha(a).$$

Set

$$(4) \quad n_\alpha(a) := x_\alpha(a)x_{-\alpha}(-a^{-1})x_\alpha(a).$$

Then,

$$\phi\left(\begin{bmatrix} 0 & a \\ -a^{-1} & 0 \end{bmatrix}\right) = n_\alpha(a).$$

We also have the following equalities,

$$(5) \quad n_\alpha(a) = x_{-\alpha}(-a^{-1})x_\alpha(a)x_{-\alpha}(-a^{-1}), \quad n_\alpha(ab) = h_\alpha(b) \cdot n_\alpha(a).$$

for any $a, b \in \mathbb{C}^*$.

Case 1: $\langle \lambda, \alpha \rangle > 0$.

Let k be any integer such that $1 \leq k \leq \langle \lambda, \alpha \rangle$. Consider the morphism $f_{\alpha,k,\lambda} : \mathbb{A}^1 \rightarrow \text{Gr}$ given by $a \mapsto x_\alpha(at^{-k+\langle \lambda, \alpha \rangle}) \cdot L_\lambda \in \mathbf{X}_\lambda$. Note that

$$(6) \quad f_{\alpha,k,\lambda}(a) = t^\lambda x_\alpha(at^{-k}) \cdot L_0.$$

As $k > 0$, $f_{\alpha,k,\lambda}$ defines an \mathbb{A}^1 -curve in \mathbf{X}_λ passing through L_λ when $a = 0$. In view of (5), we can write

$$(7) \quad x_\alpha(at^{-k}) = x_{-\alpha}(a^{-1}t^k)t^{-k\check{\alpha}}n_\alpha(a)x_{-\alpha}(a^{-1}t^k),$$

where $t^{-k\check{\alpha}} = h_\alpha(t^{-k})$. It follows that

$$f_{\alpha,k,\lambda}(a) = t^\lambda x_{-\alpha}(a^{-1}t^k)t^{-k\check{\alpha}} \cdot L_0 = t^{\lambda-k\check{\alpha}} x_{-\alpha}(a^{-1}t^{-k}) \cdot L_0.$$

When $a \rightarrow \infty$, $f_{\alpha,k,\lambda}(a) \rightarrow L_{\lambda-k\check{\alpha}}$. It follows that $L_{\lambda-k\check{\alpha}} \in \bar{\mathbf{X}}_\lambda$. In other words, $\lambda - k\check{\alpha} \in \Psi(\lambda)$.

Case 2: $\langle \lambda, \alpha \rangle < 0$.

Let k be any integer such that $1 \leq k \leq -\langle \lambda, \alpha \rangle - 1$. Consider the morphism $g_{\alpha,k,\lambda} : \mathbb{A}^1 \rightarrow \mathbf{Gr}$ given by $a \mapsto x_{-\alpha}(at^{-k-\langle \lambda, \alpha \rangle}) \cdot L_\lambda \in \mathbf{X}_\lambda$. Similar to (6), we have

$$(8) \quad g_{\alpha,k,\lambda}(a) = t^\lambda x_{-\alpha}(at^{-k}) \cdot L_0.$$

Thus, $g_{\alpha,k,\lambda}$ defines an \mathbb{A}^1 -curve in \mathbf{X}_λ passing through L_λ when $a = 0$. In view of (4), we have

$$(9) \quad x_{-\alpha}(at^{-k}) = x_\alpha(a^{-1}t^k)t^{k\check{\alpha}}n_\alpha(-a^{-1})x_{-\alpha}(a^{-1}t^k).$$

It follows that

$$g_{\alpha,k,\lambda}(a) = t^\lambda x_\alpha(a^{-1}t^k)t^{k\check{\alpha}} \cdot L_0 = t^{\lambda+k\check{\alpha}} x_\alpha(a^{-1}t^{-k}) \cdot L_0.$$

When $a \rightarrow \infty$, $g_{\alpha,k,\lambda}(a) \rightarrow L_{\lambda+k\check{\alpha}}$. It follows that $\lambda + k\check{\alpha} \in \Psi(\lambda)$. □

This lemma will provide an algorithm which completely describes the set $\Psi(\lambda)$. For a positive root α , the algorithm will rely on the sign of $\langle \lambda, \alpha \rangle$. In fact a representation-theoretic explanation will be given in Lemma 4.7 of Section 3.2.

Remark 2.3. *An analogue of Lemma 2.2 for simple roots and the highest root in the setting of Bruhat order $<$ is proved in (Io1, Lemma 1.6). In Lemma 2.22, we show that the order $<_I$ is equivalent to the Bruhat order $<$.*

For any coweight $\mu \in \check{P}$ and any positive root $\alpha \in \Phi^+$, we first introduce the following set of coweights attached to μ and α

$$(10) \quad S(\mu, \alpha) = \begin{cases} \{\mu - k\check{\alpha} \mid 0 \leq k \leq \langle \mu, \alpha \rangle\}, & \text{when } \langle \mu, \alpha \rangle \geq 0 \\ \{\mu + k\check{\alpha} \mid 0 \leq k < -\langle \mu, \alpha \rangle\}, & \text{when } \langle \mu, \alpha \rangle < 0 \end{cases}.$$

By Lemma 2.2, $S(\mu, \alpha)$ is a subset of $\Psi(\lambda)$ if $\mu \in \Psi(\lambda)$. We now define an increasing filtration $\{\Psi_i(\lambda)\}_{i \geq 0}$ of subsets in $\Psi(\lambda)$ as follows.

Definition 2.4. Define $\Psi_0(\lambda) = \{\lambda\}$, and

$$\Psi_i(\lambda) = \bigcup_{\mu \in \Psi_{i-1}(\lambda), \alpha \in \Phi^+} S(\mu, \alpha).$$

Let $\Psi_\infty(\lambda)$ denote the union of all $\Psi_i(\lambda)$. From this definition, we observe that if $\Psi_n(\lambda) = \Psi_{n+1}(\lambda)$ for some integer n , then $\Psi_\infty(\lambda) = \Psi_n(\lambda)$.

Lemma 2.5. *This filtration stabilizes after finite many steps, that is, there exists a positive integer n such that $\Psi_\infty(\lambda) = \Psi_n(\lambda)$.*

Proof. Let w be an element in the Weyl group W such that $\lambda^+ := w(\lambda)$ is dominant. Since $X_\lambda \subset \text{Gr}_{\lambda^+}$, we have

$$\Psi(\lambda) \subset \{\mu \mid L_\mu \in (\text{Gr}_{\lambda^+})^T\}.$$

Hence, $\Psi(\lambda)$ is a finite set, and so is $\Psi_\infty(\lambda)$. Therefore the filtration stabilizes after finite many steps. \square

We are now ready to state the following theorem.

Theorem 2.6. *For any $\lambda \in \check{P}$, $\Psi(\lambda) = \Psi_\infty(\lambda)$.*

This theorem gives an effective algorithm to describe the set $\Psi(\lambda)$. We will first make some preparations, and then the proof will be given in Section 2.5.

2.3. Affine Weyl group. The Weyl group W acts on the coroot lattice \check{Q} . From here we get an affine Weyl group $W_{\text{aff}} := \check{Q} \rtimes W$. We write elements of W_{aff} as $\tau_\lambda w$, where $\lambda \in \check{Q}$, $w \in W$. The element $\tau_\lambda w$ acts on \check{P} by

$$\tau_\lambda w(\mu) := w(\mu) + \lambda, \quad \text{for any } \mu \in \check{P}.$$

For any two elements $\tau_{\lambda_1} w_1, \tau_{\lambda_2} w_2$, the multiplication is given by

$$(11) \quad (\tau_{\lambda_1} w_1) \cdot (\tau_{\lambda_2} w_2) = \tau_{\lambda_1 + w_1(\lambda_2)} w_1 w_2.$$

The pair $(W_{\text{aff}}, \hat{S})$ is a Coxeter system where \hat{S} consists of simple reflections $\{s_i \mid i \in I\}$ and a simple affine reflection $s_0 = \tau_{\check{\theta}} s_\theta$ where θ is the highest positive root of G and $\check{\theta}$ is the coroot of θ . We denote by $<$ the Bruhat order on $(W_{\text{aff}}, \hat{S})$.

Let \tilde{W}_{aff} be the extended affine Weyl group $\check{P} \rtimes W$. The multiplication is given similarly as in (11). Following (IM) we define the length function ℓ on \tilde{W}_{aff} ,

$$(12) \quad \ell(\tau_\lambda w) = \sum_{\alpha \in \Phi^+, w^{-1}(\alpha) \in \Phi^+} |\langle \lambda, \alpha \rangle| + \sum_{\alpha \in \Phi^+, w^{-1}(\alpha) \in \Phi^-} |\langle \lambda, \alpha \rangle - 1|.$$

This length function ℓ on W_{aff} coincides with the length function on the Coxeter system $(W_{\text{aff}}, \hat{S})$.

Let \mathfrak{g} be the Lie algebra of G . We associate to \mathfrak{g} the (completed) affine Kac-Moody algebra $\tilde{\mathfrak{g}} := \mathfrak{g} \otimes \mathcal{K} \oplus \mathbb{C}c \oplus \mathbb{C}d$, where c is the center, d is the degree operator, and the Lie bracket is defined as in (Ka, §7.2). The affine Kac-Moody algebra $\tilde{\mathfrak{g}}$ corresponds to the extended Dynkin diagram $\hat{\Gamma}$ of \mathfrak{g} with the set of vertices $\hat{I} = I \sqcup \{0\}$. The Cartan subalgebra $\tilde{\mathfrak{h}}$ of $\tilde{\mathfrak{g}}$ is given by $\mathfrak{h} \oplus \mathbb{C}c \oplus \mathbb{C}d$, where \mathfrak{h} is the Cartan subalgebra of \mathfrak{g} . Let $\tilde{\mathfrak{h}}^*$ denote the linear dual of $\tilde{\mathfrak{h}}$. Let δ denote the linear functional on $\tilde{\mathfrak{h}}$ such that

$$\delta|_{\mathfrak{h}} = 0, \quad \delta(c) = 0, \quad \delta(d) = 1.$$

We first describe the affine root system associated to $\tilde{\mathfrak{g}}$. The set of all real affine roots of $\tilde{\mathfrak{g}}$ is given by

$$\hat{\Phi}_{\text{re}} = \{\alpha + k\delta \mid \alpha \in \Phi, k \in \mathbb{Z}\},$$

where the set of positive affine roots is given by

$$\hat{\Phi}_{\text{re}}^+ = \{\alpha + k\delta \mid \alpha \in \Phi, k > 0\} \cup \Phi^+,$$

and the set of negative affine roots is given by

$$\hat{\Phi}_{\text{re}}^- = \{\alpha + k\delta \mid \alpha \in \Phi, k < 0\} \cup \Phi^-.$$

Let $(\cdot|\cdot)$ denote the normalized bilinear form on $\tilde{\mathfrak{h}}$, and the induced bilinear form on $\tilde{\mathfrak{h}}^*$ (cf. (Ka, §6.1)). Let $\nu : \tilde{\mathfrak{h}} \rightarrow \tilde{\mathfrak{h}}^*$ be the induced isomorphism. Then $\nu(c) = \delta$. Moreover, for any $\alpha \in \Phi$, $\nu(\check{\alpha}) = \frac{2\alpha}{(\alpha|\alpha)}$ where $\check{\alpha}$ is the coroot of α . We will denote by \langle, \rangle the natural pairing between $\tilde{\mathfrak{h}}$ and $\tilde{\mathfrak{h}}^*$.

We may realize the affine Weyl group W_{aff} as the Weyl group of the affine Kac-Moody algebra $\tilde{\mathfrak{g}}$ in the sense of (Ka, §3.7), via the action of W_{aff} on $\tilde{\mathfrak{h}}^*$. Following (Ku, §13.1), we define

$$(13) \quad \tau_{\lambda} w(x) = w(x) + \langle w(x), c \rangle \nu(\lambda) - (\langle w(x), \lambda \rangle + \frac{1}{2}(\lambda|\lambda)\langle w(x), c \rangle)\delta,$$

for any $\tau_{\lambda} w \in W_{\text{aff}}$, $x \in \tilde{\mathfrak{h}}^*$.

Lemma 2.7. *The element $\tau_{k\check{\alpha}} s_{\alpha} \in W_{\text{aff}}$ corresponds to the reflection on $\tilde{\mathfrak{h}}^*$ associated to the affine root $-\alpha + k\delta$.*

Proof. For any $x \in \tilde{\mathfrak{h}}^*$, we have

$$\begin{aligned} \tau_{k\check{\alpha}} s_{\alpha}(x) &= x - \langle x, \check{\alpha} \rangle \alpha + \langle s_{\alpha}(x), c \rangle \nu(k\check{\alpha}) - (\langle s_{\alpha}(x), k\check{\alpha} \rangle + \frac{1}{2}(k\check{\alpha}|k\check{\alpha})\langle s_{\alpha}(x), c \rangle)\delta. \\ &= x - \langle x, \check{\alpha} \rangle \alpha + \frac{2k\langle x, c \rangle}{(\alpha|\alpha)} \alpha - (-k\langle x, \check{\alpha} \rangle + \frac{2k^2}{(\alpha|\alpha)}\langle x, c \rangle)\delta. \\ &= x - (-\langle x, \check{\alpha} \rangle + \frac{2k\langle x, c \rangle}{(\alpha|\alpha)})(-\alpha + k\delta) \\ &= x - \frac{2(x|-\alpha + k\delta)}{(-\alpha + k\delta|-\alpha + k\delta)}(-\alpha + k\delta), \end{aligned}$$

where the last equality holds since $\nu(c) = \delta$ and $(-\alpha + k\delta|-\alpha + k\delta) = (\alpha|\alpha)$. This is exactly the reflection on $\tilde{\mathfrak{h}}^*$ associated to $-\alpha + k\delta$. \square

For any $\alpha \in \Phi^+$ and $k \in \mathbb{Z}$, set

$$s_{\alpha, k} := \tau_{-k\check{\alpha}} s_{\alpha} \in W_{\text{aff}}.$$

Then by Lemma 2.7, $s_{\alpha, k}$ is the reflection associated to $\alpha + k\delta$. In particular $s_0 = \tau_{\theta} s_{\theta}$ is the reflection associated to the affine simple root $\alpha_0 := -\theta + \delta$.

Lemma 2.8. *Let α be a root in Φ . Assume that $k \geq 0$ if $\alpha \in \Phi^+$, and $k > 0$ if $\alpha \in \Phi^-$. Then for any $\tau_{-\lambda} w \in W_{\text{aff}}$, $s_{\alpha, k} \tau_{-\lambda} w < \tau_{-\lambda} w$ if and only if*

$$\begin{cases} k < \langle \lambda, \alpha \rangle, & \text{when } w^{-1}(\alpha) \in \Phi^+ \\ k \leq \langle \lambda, \alpha \rangle, & \text{when } w^{-1}(\alpha) \in \Phi^- \end{cases}.$$

Proof. We may realize W_{aff} as the Weyl group of the affine root system of $\tilde{\mathfrak{g}}$. Note that the assumption on α and k is equivalent to that the affine root $\alpha + k\delta$ is positive. In view of Lemma 2.7, $s_{\alpha, k}$ corresponds to the reflection $s_{\alpha + k\delta}$. By a general fact of the theory of Coxeter groups (cf. (Hu2, §5.7)), $s_{\alpha, k} \tau_{-\lambda} w < \tau_{-\lambda} w$ if and only if $w^{-1} \tau_{\lambda}(\alpha + k\delta) \in \hat{\Phi}_{\text{re}}^-$. By the formula (13),

$$w^{-1} \tau_{\lambda}(\alpha + k\delta) = w^{-1}(\alpha) + (k - \langle \lambda, \alpha \rangle)\delta.$$

Thus the lemma immediately follows. \square

Proposition 2.9. *Let α be a positive root in Φ^+ . Assume that $s_{\alpha,k}\tau_{-\lambda}w < \tau_{-\lambda}w$.*

- (1) *If $w^{-1}(\alpha) \in \Phi^+$ and $k \geq 0$, then $\langle \lambda, \alpha \rangle > 0$, and $k < \langle \lambda, \alpha \rangle$.*
- (2) *If $w^{-1}(\alpha) \in \Phi^-$ and $k \geq 0$, then $\langle \lambda, \alpha \rangle \geq 0$, and $k \leq \langle \lambda, \alpha \rangle$.*
- (3) *If $w^{-1}(\alpha) \in \Phi^+$ and $k < 0$, then $\langle \lambda, \alpha \rangle < 0$, and $k \geq \langle \lambda, \alpha \rangle$.*
- (4) *If $w^{-1}(\alpha) \in \Phi^-$ and $k < 0$, then $\langle \lambda, \alpha \rangle < -1$, and $k > \langle \lambda, \alpha \rangle$.*

Remark 2.10. *When α is a simple root or a highest root, a similar characterization appears in (Io1, Lemma 1.3).*

Proof. In first two cases, $s_{\alpha,k}$ corresponds to the reflection of the positive affine root $\alpha + k\delta$. In the last two cases, $s_{\alpha,k}$ corresponds to the reflection of the positive affine root $-\alpha - k\delta$. Then the proposition easily follows from Lemma 2.8. \square

Definition 2.11. A coweight $\check{\omega} \in \check{P}$ is called miniscule, if for any positive root $\alpha \in \Phi^+$, $\langle \check{\omega}, \alpha \rangle \in \{0, 1\}$.

The following corollary and Proposition 2.9 will be used in Section 2.5.

Corollary 2.12. *Let $\check{\omega}$ be a miniscule coweight. For any $\lambda \in \check{Q} - \check{\omega}$, $\alpha \in \Phi^+$ and $y \in \tau_{\check{\omega}}W\tau_{-\check{\omega}}$, assume that $s_{\alpha,k}\tau_{-\lambda-\check{\omega}}y < \tau_{-\lambda-\check{\omega}}y$.*

- (1) *If $k \geq 0$, then $k \leq \langle \lambda, \alpha \rangle$;*
- (2) *If $k < 0$, then $k \geq \langle \lambda, \alpha \rangle$.*

Proof. We can write $y = \tau_{\check{\omega}}w\tau_{-\check{\omega}}$ for some $w \in W$. Note that

$$\tau_{-\lambda-\check{\omega}}y = \tau_{-(\lambda+w(\check{\omega}))}w.$$

We first prove part (1). Assume that $k \geq 0$. If $w^{-1}(\alpha) \in \Phi^+$, then $\langle w(\check{\omega}), \alpha \rangle = \langle \check{\omega}, w^{-1}(\alpha) \rangle \in \{0, 1\}$. Moreover, by part (1) of Proposition 2.9,

$$k < \langle \lambda + w(\check{\omega}), \alpha \rangle = \langle \lambda, \alpha \rangle + \langle w(\check{\omega}), \alpha \rangle.$$

Hence $k \leq \langle \lambda, \alpha \rangle$. If $w^{-1}(\alpha) \in \Phi^-$, then $\langle w(\check{\omega}), \alpha \rangle \in \{0, -1\}$. By part (2) of Proposition 2.9, we get

$$k \leq \langle \lambda + w(\check{\omega}), \alpha \rangle = \langle \lambda, \alpha \rangle + \langle w(\check{\omega}), \alpha \rangle.$$

Hence we also have $k \leq \langle \lambda, \alpha \rangle$. This shows that in case $k \geq 0$, we always have $k \leq \langle \lambda, \alpha \rangle$, no matter $w^{-1}(\alpha)$ is positive or negative.

By similar arguments, we can show that if $k < 0$, then $k \geq \langle \lambda, \alpha \rangle$. \square

2.4. Components of Gr as partial flag varieties of the Kac-Moody group. Let G_{sc} be the simply-connected cover of G . Let \tilde{G}_{sc} be the affine Kac-Moody group with Lie algebra $\tilde{\mathfrak{g}}$ in the sense of Kumar and Mathieu (cf. (Ku, §VI)), which can be realized as a central extension of the semi-direct product $G_{\text{sc}}(\mathcal{K}) \rtimes \mathbb{C}^\times$ (\mathbb{C}^\times acts on $G_{\text{sc}}(\mathcal{K})$ by the loop rotation) (cf. (Ku, Theorem 13.2.8)). It is known that when G is of adjoint type, the associated affine Grassmannian Gr has $|\check{P}/\check{Q}|$ components. In this subsection we produce an explicit description of each component of Gr as a partial flag variety of the Kac-Moody group \tilde{G}_{sc} .

Let \hat{M} denote the set of vertices $i \in \hat{I}$ such that $a_i = 1$, where a_i is the Kac labeling of affine Dynkin diagram $\hat{\Gamma}$ (Ka, p.54, Table Aff 1). Since $a_0 = 1$, $\hat{M} = M \cup \{0\}$, where $M := \hat{M} \cap I$. Let θ be the highest root of \mathfrak{g} . It is known that

$$(14) \quad \theta = \sum_{i \in I} a_i \alpha_i.$$

For each $i \in I$, let $\check{\omega}_i$ be the fundamental coweight of \mathfrak{g} attached to i , i.e. for any simple root α_j ,

$$\langle \check{\omega}_i, \alpha_j \rangle = \delta_{i,j}.$$

Lemma 2.13. *For any $i \in I$, the fundamental coweight $\check{\omega}_i$ is miniscule (cf. Definition 2.11) if and only if $i \in M$, if and only if $\langle \check{\omega}_i, \theta \rangle = 1$.*

Proof. This can be read from the Kac labeling in (Ka, p.54, Table Aff 1) for $\tilde{\mathfrak{g}}$, and the list of miniscule fundamental weights for the dual Lie algebra ${}^L\mathfrak{g}$ of \mathfrak{g} in (Hu1, p.72, ex.13) \square

Lemma 2.14. *The set $\{\check{\omega}_\kappa \mid \kappa \in \hat{M}\}$ gives a complete set of coset representatives of \check{P}/\check{Q} , as does the set $\{-\check{\omega}_\kappa \mid \kappa \in \hat{M}\}$.*

Proof. This appears as Corollary to Prop. 6 in (Bou, Chapter VI.2.3). \square

For any coweight $\mu \in \check{P}$, we define a conjugation $\text{Ad}_\mu : W_{\text{aff}} \rightarrow W_{\text{aff}}$ given by

$$\text{Ad}_\mu(\tau_{\lambda w}) := \tau_\mu(\tau_{\lambda w})\tau_{-\mu}, \text{ for any } \tau_{\lambda w} \in W_{\text{aff}}.$$

For each $\kappa \in \hat{M}$, set $W_\kappa = \text{Ad}_{\check{\omega}_\kappa}(W)$, where by convention $\check{\omega}_0 = 0$. Set $I_\kappa = \hat{I} \setminus \{\kappa\}$. The following proposition is well-known.

For the convenience of the reader, we provide a proof here.

Proposition 2.15. *For any $\kappa \in \hat{M}$, the subgroup W_κ is a parabolic subgroup of W_{aff} with Coxeter generators $\{s_i \mid i \in I_\kappa\}$.*

Proof. When $\kappa = 0$, $W_\kappa = W$ together with $\{s_i \mid i \in I\}$ is clearly a Coxeter system.

Now we assume that $\kappa \in M$. For any $i \in \hat{I} \setminus \{0, \kappa\}$, it is easy to see that $\text{Ad}_{\check{\omega}_\kappa}(s_i) = s_i$. Since $\theta = \sum_{i \in I} a_i \alpha_i$, by Lemma 2.13 we have

$$\text{Ad}_{\check{\omega}_\kappa}(s_\theta) = \tau_{\check{\theta}} s_\theta = s_0.$$

Thus, W_κ contains $\{s_i \mid i \in I_\kappa\}$. Let W'_κ be the subgroup of W_κ generated by $\{s_i \mid i \in I_\kappa\}$. As we see in the table (Ka, p.54, Table Aff 1), by deleting κ the Dynkin diagram $\hat{\Gamma} \setminus \{\kappa\}$ is the same as the Dynkin diagram Γ of \mathfrak{g} . Therefore W'_κ isomorphic to W as Coxeter groups, in particular $|W'_\kappa| = |W|$. On the other hand the conjugation $\text{Ad}_{\check{\omega}_\kappa}$ also gives rise to an isomorphism $W \simeq W_\kappa$ of finite groups. It follows that $|W_\kappa| = |W'_\kappa|$. Therefore $W_\kappa = W'_\kappa$, and furthermore W_κ is a parabolic subgroup of W_{aff} with Coxeter generators $\{s_i \mid i \in I_\kappa\}$. \square

From this proposition, we may deduce an interesting corollary on the finite Weyl group W .

Corollary 2.16. *The Weyl group W can be generated by the set of reflections $\{s_i \mid i \in I \setminus \{\kappa\}\} \cup \{s_\theta\}$ for any $\kappa \in M$ (equivalently, the coefficient of α_κ in the highest root θ is 1).*

Proof. By Proposition 2.15, the set $\{\text{Ad}_{-\check{\omega}_k}(s_i) \mid i \in I_k\}$ generates the Weyl group W . Unfolding the elements $\text{Ad}_{-\check{\omega}_k}(s_i)$, the corollary immediately follows. \square

The set $\pi_0(\text{Gr})$ of the components of the affine Grassmannian Gr can be identified with \check{P}/\check{Q} . Let Gr^κ be the component of Gr containing the point $L_{-\check{\omega}_k} = t^{-\check{\omega}_k}G(\mathcal{O})/G(\mathcal{O})$. By Lemma 2.14, we have the following disjoint union decomposition of Gr

$$\text{Gr} = \sqcup_{\kappa \in \hat{M}} \text{Gr}^\kappa.$$

For any $\kappa \in \hat{M}$, set

$$(15) \quad \check{Q}_\kappa := \check{Q} - \check{\omega}_\kappa.$$

Any T -fixed point in Gr^κ is given by L_λ and any I -orbit in Gr^κ is given by X_λ , for some $\lambda \in \check{Q}_\kappa$. Note that there is a bijection $\iota_\kappa : \check{Q}_\kappa \rightarrow W_{\text{aff}}/W_\kappa$ given by

$$\lambda \mapsto \tau_{-\lambda - \check{\omega}_\kappa} W_\kappa,$$

which fits into the following commutative diagram

$$\begin{array}{ccc} \check{Q} & \xrightarrow{\iota_0} & W_{\text{aff}}/W \\ \cdot \check{\omega}_\kappa \downarrow & & \downarrow \text{Ad}_{\check{\omega}_\kappa} \\ \check{Q}_\kappa & \xrightarrow{\iota_\kappa} & W_{\text{aff}}/W_\kappa \end{array}.$$

In the following we would like to describe the component Gr^κ as a partial flag variety associated to the quotient W_{aff}/W_κ .

There is a canonical projection $\pi : \tilde{G}_{\text{sc}} \rightarrow G_{\text{sc}}(\mathcal{H}) \times \mathbb{C}^\times$. The preimage

$$\tilde{B} = \pi^{-1}(\mathbb{I}_{\text{sc}} \times \mathbb{C}^\times)$$

is the standard Borel subgroup of \tilde{G}_{sc} , where \mathbb{I}_{sc} is the Iwahori subgroup in $G_{\text{sc}}(\mathcal{H})$. We have the following identification

$$\tilde{G}_{\text{sc}}/\tilde{B} \simeq G_{\text{sc}}(\mathcal{H})/\mathbb{I}_{\text{sc}}.$$

For any $\tau_\lambda w \in W_{\text{aff}}$, we associate an \mathbb{I}_{sc} -orbit

$$Y_{\tau_\lambda w} := \mathbb{I}_{\text{sc}} t^{-\lambda} w \mathbb{I}_{\text{sc}} / \mathbb{I}_{\text{sc}} \subset G_{\text{sc}}(\mathcal{H})/\mathbb{I}_{\text{sc}}.$$

Then $\dim Y_{\tau_\lambda w} = \ell(\tau_\lambda w)$.

Remark 2.17. *The sign normalization in $Y_{\tau_\lambda w}$ is crucial. Without this sign, the dimension formula for $\dim Y_{\tau_\lambda w}$ does not hold.*

Let \tilde{P}_κ denote the maximal standard parabolic subgroup of \tilde{G}_{sc} containing \tilde{B} , which is associated to the subset $I_\kappa = \hat{I} \setminus \{\kappa\}$ of \hat{I} . By Proposition 2.15, \tilde{P}_κ corresponds to the parabolic subgroup W_κ of W_{aff} . The \tilde{B} -orbits in the partial flag variety $\tilde{G}_{\text{sc}}/\tilde{P}_\kappa$ can be indexed by the cosets in W_{aff}/W_κ .

Without confusion, we still denote by $p : G_{\text{sc}}(\bar{\mathcal{H}}) \rightarrow G(\bar{\mathcal{H}})$ and $p : G_{\text{sc}}(\mathcal{H}) \rightarrow G(\mathcal{H})$ the maps induced from the covering map $p : G_{\text{sc}} \rightarrow G$, where $\bar{\mathcal{H}}$ is the algebraic closure of \mathcal{H} . The map $p : G_{\text{sc}}(\bar{\mathcal{H}}) \rightarrow G(\bar{\mathcal{H}})$ is surjective, but $p : G_{\text{sc}}(\mathcal{H}) \rightarrow G(\mathcal{H})$ is not if $\check{P}/\check{Q} \neq 0$.

Let \tilde{t}_κ be a lifting of $t^{-\check{\omega}_\kappa} \in G(\mathcal{H})$ in $G_{\text{sc}}(\tilde{\mathcal{H}})$ via the map p . Let $\text{Ad}_{\tilde{t}_\kappa}$ denote the conjugation map on $G_{\text{sc}}(\tilde{\mathcal{H}})$ by \tilde{t}_κ , i.e.

$$\text{Ad}_{\tilde{t}_\kappa}(g) := \tilde{t}_\kappa g \tilde{t}_\kappa^{-1}, \text{ for any } g \in G_{\text{sc}}(\tilde{\mathcal{H}}).$$

Lemma 2.18. *The conjugation $\text{Ad}_{\tilde{t}_\kappa}$ preserves $G_{\text{sc}}(\mathcal{H})$, and $\text{Ad}_{\tilde{t}_\kappa}$ is independent of the choice of the lifting \tilde{t}_κ .*

Proof. For any root $\alpha \in \Phi$, we have

$$(16) \quad \tilde{t}_\kappa x_\alpha(a) \tilde{t}_\kappa^{-1} = x_\alpha(at^{-\langle \check{\omega}_\kappa, \alpha \rangle}),$$

where x_α is a root subgroup homomorphism of G_{sc} associated to α , and $a \in \mathcal{H}$. since the group $G_{\text{sc}}(\mathcal{H})$ is generated by its root subgroups $x_{\alpha_i}(\mathcal{H})$ (cf. (Stg, §7, Theorem 10)), it follows that the conjugation $\text{Ad}_{\tilde{t}_\kappa}$ preserves $G_{\text{sc}}(\mathcal{H})$. The independence of the lifting \tilde{t}_κ also follows from the formula (16). \square

Set

$$G_{\text{sc}}(\mathcal{O})_\kappa := \text{Ad}_{\tilde{t}_\kappa}(G_{\text{sc}}(\mathcal{O})).$$

Lemma 2.19. *For any $\kappa \in \hat{M}$, the parabolic subgroup \tilde{P}_κ is equal to the preimage $\pi^{-1}(G_{\text{sc}}(\mathcal{O})_\kappa \rtimes \mathbb{C}^\times)$, where π is the canonical projection $\pi : \tilde{G}_{\text{sc}} \rightarrow G_{\text{sc}}(\mathcal{H}) \rtimes \mathbb{C}^\times$.*

Proof. We choose root subgroup homomorphisms $\hat{x}_{\alpha_i}, \hat{x}_{-\alpha_i} : \mathbb{C} \rightarrow \tilde{G}_{\text{sc}}$ associated to α_i for each $i \in \hat{I}$. For each $i \in I$, set $x_{\alpha_i} := \pi \circ \hat{x}_{\alpha_i}$ and $x_{-\alpha_i} := \pi \circ \hat{x}_{-\alpha_i}$. The images of $x_{\pm\alpha_i}$ land in G_{sc} , and they are exactly the root subgroup homomorphisms of G_{sc} associated to $\pm\alpha_i$ for each $i \in I$. When $i = 0$, there exist root subgroup homomorphisms $x_\theta, x_{-\theta}$ of G_{sc} associated to $\theta, -\theta$, such that

$$\pi \circ \hat{x}_{\alpha_0}(a) = x_{-\theta}(at), \quad \pi \circ \hat{x}_{-\alpha_0}(a) = x_\theta(at^{-1}), \quad \text{for any } a \in \mathbb{C}.$$

Set $\hat{P}'_\kappa := \pi^{-1}(G_{\text{sc}}(\mathcal{O})_\kappa \rtimes \mathbb{C}^\times)$. We first observe that $\tilde{B} \subset \hat{P}'_\kappa$, since $\langle \check{\omega}_\kappa, \alpha \rangle \in \{0, 1\}$ for any $\kappa \in \hat{M}$ and any positive root $\alpha \in \Phi^+$. Hence \hat{P}'_κ is a standard parabolic subgroup of \tilde{G}_{sc} (cf. (Ku, Theorem 5.1.3, Theorem 6.1.17)). To show $\hat{P}'_\kappa = \tilde{P}_\kappa$, it suffices to check that \hat{P}'_κ corresponds to the subset $I_\kappa \subset \hat{I}$. In other words, it suffices to check that \hat{P}'_κ is proper and contains all root subgroups $\hat{x}_{-\alpha_i}$ for all $i \in I_\kappa$. Equivalently it suffices to show that $G(\mathcal{O})_\kappa$ contains $\pi \circ \hat{x}_{-\alpha_i}$ for all $i \in I_\kappa$.

When $\kappa = 0$, it suffices to check that $G(\mathcal{O})$ contains $x_{-\alpha_i}$ for each $i \in I$, as well as $x_{-\theta}(\cdot t)$. This is obvious. When $\kappa \in M$, it suffices to show that $G_{\text{sc}}(\mathcal{O})_\kappa$ is a proper subgroup of $G_{\text{sc}}(\mathcal{H})$, and $G_{\text{sc}}(\mathcal{O})_\kappa$ contains $x_{-\alpha_i}$ for each $i \in I \setminus \{\kappa\}$, and $x_\theta(\cdot t^{-1})$. Clearly $G_{\text{sc}}(\mathcal{O})_\kappa$ is a proper subgroup of $G_{\text{sc}}(\mathcal{H})$, since $G_{\text{sc}}(\mathcal{O})$ is proper in $G_{\text{sc}}(\mathcal{H})$ and by Lemma 2.18 $\text{Ad}_{\tilde{t}_\kappa}$ preserves $G_{\text{sc}}(\mathcal{H})$. The group $G_{\text{sc}}(\mathcal{O})_\kappa$ contains $x_{-\alpha_i}$ for each $i \in I \setminus \{\kappa\}$, since

$$\tilde{t}_\kappa x_{-\alpha_i}(a) \tilde{t}_\kappa^{-1} = x_{-\alpha_i}(a).$$

Recall that $\langle \check{\omega}_\kappa, \theta \rangle = 1$. By the computation

$$\tilde{t}_\kappa x_\theta(a) \tilde{t}_\kappa^{-1} = x_\theta(at^{-1}),$$

we see that $G_{\text{sc}}(\mathcal{O})_\kappa$ contains $x_\theta(at^{-1})$ for any $a \in \mathbb{C}$. This completes the proof. \square

Example 2.20. We examine $G = \mathrm{PGL}_3$. In this case $G_{\mathrm{sc}} = \mathrm{SL}_3$. All fundamental coweights are miniscule. Let $\check{\omega}_1$ be the first fundamental coweight. Then

$$t^{-\check{\omega}_1} = \begin{pmatrix} t^{-1} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \in \mathrm{PGL}_3(\mathcal{K})$$

has a lifting

$$\tilde{t}^{-\check{\omega}_1} = \begin{pmatrix} t^{-2/3} & 0 & 0 \\ 0 & t^{1/3} & 0 \\ 0 & 0 & t^{1/3} \end{pmatrix} \in \mathrm{SL}_3(\tilde{\mathcal{K}}).$$

By conjugating we see

$$\mathrm{SL}_3(\mathcal{O})_1 := \mathrm{Ad}_{\tilde{t}^{-\check{\omega}_1}}(\mathrm{SL}_3(\mathcal{O})) = \left\{ \begin{pmatrix} a_{11} & t^{-1}a_{12} & t^{-1}a_{12} \\ ta_{21} & a_{22} & a_{23} \\ ta_{31} & a_{32} & a_{33} \end{pmatrix} \middle| (a_{ij}) \in \mathrm{SL}_3(\mathcal{O}) \right\}.$$

The group $p^{-1}(\mathrm{SL}_3(\mathcal{O})_1 \rtimes \mathbb{C}^\times)$ is the maximal parabolic subgroup of the Kac-Moody group $\widetilde{\mathrm{SL}}_3$ associated to the set of simple roots $\{\alpha_0, \alpha_2\}$.

Proposition 2.21. *There exists an isomorphism $i_\kappa : \mathrm{Gr}^\kappa \rightarrow \tilde{G}_{\mathrm{sc}}/\tilde{P}_\kappa$ as \tilde{G}_{sc} -homogeneous spaces with $i_\kappa(L_{-\check{\omega}_\kappa}) = e\tilde{P}_\kappa$, and moreover, $i_\kappa(X_\lambda) = \tilde{B}\tau_{-\lambda-\check{\omega}_\kappa}\tilde{P}_\kappa/\tilde{P}_\kappa$.*

Proof. It is well-known that the central component Gr^0 is isomorphic to $G_{\mathrm{sc}}(\mathcal{K})/G_{\mathrm{sc}}(\mathcal{O})$ as a homogenous space (this can be seen by comparing the Cartan decompositions for $G(\mathcal{K})$ and $G_{\mathrm{sc}}(\mathcal{K})$).

The translation map from Gr^0 to Gr^κ given by $L \mapsto t^{-\check{\omega}_\kappa}L$ is an isomorphism between Gr^0 and Gr^κ . In view of Lemma 2.18, one can see that the component Gr^κ is also a homogeneous space of $G_{\mathrm{sc}}(\mathcal{K})$. Moreover $G_{\mathrm{sc}}(\mathcal{O})_\kappa$ is exactly the stabilizer group of $G_{\mathrm{sc}}(\mathcal{K})$ at $L_{-\check{\omega}_\kappa} \in \mathrm{Gr}^\kappa$ (recall that \tilde{t}_κ is a lifting of $t^{-\check{\omega}_\kappa}$). Hence $\mathrm{Gr}^\kappa \simeq G_{\mathrm{sc}}(\mathcal{K})/G_{\mathrm{sc}}(\mathcal{O})_\kappa$. Furthermore

$$\mathrm{Gr}^\kappa \simeq (G_{\mathrm{sc}}(\mathcal{K}) \rtimes \mathbb{C}^\times)/(G_{\mathrm{sc}}(\mathcal{O})_\kappa \rtimes \mathbb{C}^\times),$$

since the rotation factor \mathbb{C}^\times acts trivially at $L_{-\check{\omega}_\kappa}$.

Finally we consider the action of \tilde{G}_{sc} on Gr^κ , which factors through the action of $G_{\mathrm{sc}}(\mathcal{K}) \rtimes \mathbb{C}^\times$. The stabilizer group of \tilde{G}_{sc} at $L_{-\check{\omega}_\kappa}$ is $p^{-1}(G_{\mathrm{sc}}(\mathcal{O})_\kappa \rtimes \mathbb{C}^\times)$. By Lemma 2.19, we conclude that $\mathrm{Gr}^\kappa \simeq \tilde{G}_{\mathrm{sc}}/\tilde{P}_\kappa$.

For any $\lambda \in \check{Q}_\kappa$, equivalently $\lambda + \check{\omega}_\kappa \in \check{Q}$, so we have $t^{\lambda+\check{\omega}_\kappa} \in G_{\mathrm{sc}}(\mathcal{K})$. Then

$$\tilde{B}\tau_{-\lambda-\check{\omega}_\kappa}\tilde{P}_\kappa/\tilde{P}_\kappa \simeq \mathrm{I}_{\mathrm{sc}} t^{\lambda+\check{\omega}_\kappa} \cdot L_{-\check{\omega}_\kappa} = \mathrm{I}_{\mathrm{sc}} \cdot L_\lambda = \mathrm{I} \cdot L_\lambda = X_\lambda,$$

since the natural map $\mathrm{I}_{\mathrm{sc}} \rightarrow \mathrm{I}$ is surjective. This completes the proof of this proposition. \square

Given any two coweights $\lambda, \mu \in \check{Q}_\kappa$, let $\tau_{-\lambda-\check{\omega}_\kappa}y$ (resp. $\tau_{-\mu-\check{\omega}_\kappa}w$) be the minimal representative in the coset $\tau_{-\lambda-\check{\omega}_\kappa}W_\kappa$ (resp. $\tau_{-\mu-\check{\omega}_\kappa}W_\kappa$).

Lemma 2.22. *$\lambda <_I \mu$ if and only if $\tau_{-\lambda-\check{\omega}_\kappa}y < \tau_{-\mu-\check{\omega}_\kappa}w$, where $<$ is the Bruhat order on W_{aff} .*

Proof. This equivalence follows from Proposition 2.21 and (Ku, Proposition 7.1.21), $\mu <_I \lambda$. \square

2.5. Proof of Theorem 2.6. We first recall the statement of Theorem 2.6.

Theorem. For any $\lambda \in \check{P}$, $\Psi(\lambda) = \Psi_\infty(\lambda)$.

Proof. By Lemma 2.2, it suffices to show that $\mu \in \Psi(\lambda)$ implies $\mu \in \Psi_\infty(\lambda)$. We may assume that $\lambda, \mu \in \check{Q}_\kappa$ for some $\kappa \in \hat{M}$.

Let $\tau_{-\lambda-\check{\omega}_\kappa}y$ (resp. $\tau_{-\mu-\check{\omega}_\kappa}w$) be the minimal representative in the coset $\tau_{-\lambda-\check{\omega}_\kappa}W_\kappa$ (resp. $\tau_{-\mu-\check{\omega}_\kappa}W_\kappa$). By Lemma 2.22, $\mu <_I \lambda$ if and only if $\tau_{-\mu-\check{\omega}_\kappa}y < \tau_{-\lambda-\check{\omega}_\kappa}w$.

By the chain property of partial Bruhat order for Coxeter groups (cf. (BB, Theorem 2.5.5)), there exists a sequence of elements

$$(17) \quad \tau_{-\mu-\check{\omega}_\kappa}w = \tau_{-\lambda_n-\check{\omega}_\kappa}y_n < \tau_{-\lambda_{n-1}-\check{\omega}_\kappa}y_{n-1} < \cdots < \tau_{-\lambda_1-\check{\omega}_\kappa}y_1 < \tau_{-\lambda_0-\check{\omega}_\kappa}y_0 = \tau_{-\lambda-\check{\omega}_\kappa}y,$$

which satisfies the following properties:

- (1) for each $0 \leq i \leq n$, $\tau_{-\lambda_i-\check{\omega}_\kappa}y_i$ is the minimal representative in the coset $\tau_{-\lambda_i-\check{\omega}_\kappa}W_\kappa$;
- (2) for each $1 \leq i \leq n$, $\tau_{-\lambda_i-\check{\omega}_\kappa}y_i = s_{\beta_i, k_i} \tau_{-\lambda_{i-1}-\check{\omega}_\kappa}y_{i-1}$ for some affine reflection s_{β_i, k_i} with $\beta_i \in \Phi^+$ and for some $k_i \in \mathbb{Z}$, and $\ell(\tau_{-\lambda_i-\check{\omega}_\kappa}y_i) = \ell(\tau_{-\lambda_{i-1}-\check{\omega}_\kappa}y_{i-1}) - 1$.

By the choice of this sequence, we see that all λ_i are distinct. By Proposition 2.21 and (Ku, Proposition 7.1.21) again, the sequence (17) is equivalent to

$$\lambda_n = \mu <_I \lambda_{n-1} <_I \cdots <_I \lambda_i \cdots <_I \lambda_1 <_I \lambda_0 = \lambda.$$

From the following computation

$$\tau_{-\lambda_i-\check{\omega}_\kappa}y_i = \tau_{-k_i\check{\beta}_i} s_{\beta_i} \tau_{-\lambda_{i-1}-\check{\omega}_\kappa}y_{i-1} = \tau_{-\lambda_{i-1} + (\langle \lambda_{i-1}, \beta_i \rangle - k_i)\check{\beta}_i - \check{\omega}_\kappa} \text{Ad}_{\check{\omega}_\kappa}(s_{\beta_i})y_{i-1},$$

we see that $\lambda_i = \lambda_{i-1} - (\langle \lambda_{i-1}, \beta_i \rangle - k_i)\check{\beta}_i$ and $y_i = \text{Ad}_{\check{\omega}_\kappa}(s_{\beta_i})y_{i-1} \in W_\kappa$. It follows that $\langle \lambda_{i-1}, \beta_i \rangle - k_i \neq 0$, since all λ_i are distinct. Since

$$\tau_{-\lambda_{i-1}-\check{\omega}_\kappa}y_{i-1} < \tau_{-\lambda_i-\check{\omega}_\kappa}y_i = s_{\beta_i, k_i} \tau_{-\lambda_{i-1}-\check{\omega}_\kappa}y_{i-1},$$

in view of Proposition 2.9 and Corollary 2.12, we have

- (1) if $k \geq 0$, then $0 < \langle \lambda_{i-1}, \beta_i \rangle - k_i \leq \langle \lambda_{i-1}, \beta_i \rangle$;
- (2) if $k < 0$, then $0 < k_i - \langle \lambda_{i-1}, \beta_i \rangle < -\langle \lambda_{i-1}, \beta_i \rangle$.

Therefore $\lambda_i \in \Psi_1(\lambda_{i-1})$ for any $1 \leq i \leq n$ (cf. Definition 2.4). In fact $\lambda_i \in \Psi_i(\lambda_0)$ for each i . It follows that $\mu \in \Psi_\infty(\lambda)$, since by convention $\mu = \lambda_n$ and $\lambda = \lambda_0$. \square

2.6. Crossing different components of Gr. Recall that \check{W}_{aff} is the extended affine Weyl group $\check{P} \rtimes W$. Let Ω denote the group of all length zero elements in \check{W}_{aff} . Equivalently, Ω is the stabilizer group of \check{W}_{aff} at the fundamental alcove of W_{aff} (cf. (Hu2, §4.5)). Then $\Omega \simeq \check{P}/\check{Q}$. Let $W_{\check{\omega}_\kappa}$ denote the stabilizer group of W at $\check{\omega}_\kappa$. Then $W_{\check{\omega}_\kappa}$ is a parabolic subgroup of W with Coxeter generators $\{s_i \mid i \in \hat{I} \setminus \{0, \kappa\}\}$. For each $\kappa \in \hat{M}$, let w_κ denote the longest element in $W_{\check{\omega}_\kappa}$ for each $\kappa \in \hat{M}$, in particular w_0 is the longest element in W . For each $\kappa \in \hat{M}$, set $w^\kappa = w_\kappa w_0$.

Lemma 2.23. The group Ω consists of elements

$$\gamma_\kappa := \tau_{\check{\omega}_\kappa} w^\kappa \in \check{W}_{\text{aff}}, \kappa \in \hat{M},$$

Proof. See (Mac, 2.5.4). \square

Using the length zero element γ_κ , we define a translation map $\rho_\kappa : \text{Gr}^0 \rightarrow \text{Gr}^\kappa$ given by

$$L \mapsto t^{-\check{\omega}_\kappa} w^\kappa L \in \text{Gr}^\kappa, \text{ for any } L \in \text{Gr}^0.$$

Proposition 2.24. *For any $\lambda \in \check{Q}$, we have $\rho_\kappa(\mathbf{X}_\lambda) = \mathbf{X}_{w^\kappa(\lambda) - \check{\omega}_\kappa}$, and $\rho_\kappa(\bar{\mathbf{X}}_\lambda) = \bar{\mathbf{X}}_{w^\kappa(\lambda) - \check{\omega}_\kappa}$.*

Proof. Since the map ρ_κ is an isomorphism between Gr^0 and Gr^κ , it suffices to show that $\rho_\kappa(\mathbf{X}_\lambda) = \mathbf{X}_{w^\kappa(\lambda) - \check{\omega}_\kappa}$. We are reduced to show the following fact

$$t^{-\check{\omega}_\kappa} w^\kappa \mathbf{I}(w^\kappa)^{-1} t^{\check{\omega}_\kappa} = \mathbf{I}.$$

For any $\alpha \in \Phi^+$,

$$t^{-\check{\omega}_\kappa} w^\kappa x_\alpha(a) (w^\kappa)^{-1} t^{\check{\omega}_\kappa} = x_{w^\kappa(\alpha)}(at^{-\langle \check{\omega}_\kappa, w^\kappa(\alpha) \rangle}),$$

where x_α (resp. $x_{w^\kappa(\alpha)}$) is the root group homomorphism associated to α (resp. $w^\kappa(\alpha)$), and $a \in \mathbb{C}$. Notice that $w^\kappa(\alpha) \in \Phi^-$ if and only if the support of $w^\kappa(\alpha)$ contains the simple root α_κ with coefficient -1 . It follows that

$$-\langle \check{\omega}_\kappa, w^\kappa(\alpha) \rangle = \begin{cases} 1 & \text{if } w^\kappa(\alpha) \in \Phi^- \\ 0 & \text{if } w^\kappa(\alpha) \in \Phi^+ \end{cases}.$$

It follows that

$$t^{-\check{\omega}_\kappa} w^\kappa x_\alpha(a) (w^\kappa)^{-1} t^{\check{\omega}_\kappa} \in \mathbf{I}.$$

For any $\alpha \in \Phi^-$,

$$t^{-\check{\omega}_\kappa} w^\kappa x_\alpha(at) (w^\kappa)^{-1} t^{\check{\omega}_\kappa} = x_{w^\kappa(\alpha)}(at^{1 - \langle \check{\omega}_\kappa, w^\kappa(\alpha) \rangle}).$$

In this case $w^\kappa(\alpha) \in \Phi^+$ if and only if the support of $w^\kappa(\alpha)$ contains the simple root α_κ with coefficient 1 . It follows that

$$1 - \langle \check{\omega}_\kappa, w^\kappa(\alpha) \rangle = \begin{cases} 0 & \text{if } w^\kappa(\alpha) \in \Phi^+ \\ 1 & \text{if } w^\kappa(\alpha) \in \Phi^- \end{cases}.$$

Therefore

$$t^{-\check{\omega}_\kappa} w^\kappa x_\alpha(at) (w^\kappa)^{-1} t^{\check{\omega}_\kappa} \in \mathbf{I}.$$

This finishes the proof. \square

Now we define a map $\bar{\rho}_\kappa : \check{Q} \rightarrow \check{Q}_\kappa$, given by $\bar{\rho}_\kappa(\lambda) = w^\kappa(\lambda) - \check{\omega}_\kappa$. The following corollary is a consequence of Proposition 2.24.

Corollary 2.25. *The map $\bar{\rho}_\kappa$ gives rise to a bijection from $\Psi(\lambda)$ to $\Psi(w^\kappa(\lambda) - \check{\omega}_\kappa)$ which preserves the partial order $<_I$, for any $\lambda \in \check{Q}$ and $\kappa \in \hat{M}$.*

Notice that for any $\alpha \in \Phi^+$, if $\langle \lambda, \alpha \rangle > 0$ and $0 \leq k \leq \langle \lambda, \alpha \rangle$, by Corollary 2.25

$$\bar{\rho}_\kappa(\lambda - k\check{\alpha}) = \bar{\rho}_\kappa(\lambda) - kw^\kappa(\check{\alpha}) \in \Psi(w^\kappa(\lambda) - \check{\omega}_\kappa).$$

Given a positive root α satisfying above conditions, $w^\kappa(\alpha)$ could be a negative root. If $w^\kappa(\alpha) \in \Phi^-$, then

$$\langle \bar{\rho}_\kappa(\lambda), -w^\kappa(\alpha) \rangle < 0.$$

In this case, $\lambda - k\check{\alpha}$ is obtained by successively subtracting the positive coroot $\check{\alpha}$, however $\bar{\rho}_\kappa(\lambda - k\alpha)$ is obtained by successively adding the positive coroot $-w^\kappa(\check{\alpha})$. There is a similar phenomenon when $\langle \lambda, \alpha \rangle < 0$.

3. A DUALITY BETWEEN AFFINE SCHUBERT VARIETIES AND LEVEL ONE AFFINE DEMAZURE MODULES

3.1. **The Kac-Moody algebra ${}^L\tilde{\mathfrak{g}}$.** Let ${}^L\hat{\Gamma}$ be the Dynkin diagram which is dual to the Dynkin diagram $\hat{\Gamma}$ of $\tilde{\mathfrak{g}}$. Let ${}^L\tilde{\mathfrak{g}}$ denote the Kac-Moody algebra associated to ${}^L\hat{\Gamma}$. Let $\{\check{e}_i, \check{f}_i | i \in \hat{I}\}$ be a set of Chevalley generators of ${}^L\tilde{\mathfrak{g}}$. Let ${}^L\mathfrak{g}$ be the Lie subalgebra of ${}^L\tilde{\mathfrak{g}}$ generated by $\{\check{e}_i, \check{f}_i | i \in I\}$. Then ${}^L\mathfrak{g}$ is a simple Lie algebra with Dynkin diagram ${}^L\Gamma$ which is dual to the Dynkin diagram Γ of \mathfrak{g} . We have the following table for the correspondence between $(\Gamma, \hat{\Gamma})$ and $({}^L\Gamma, {}^L\hat{\Gamma})$

(18)

| | | | | | | | | | |
|--------------------|-------------|------------------|-----------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Γ | A_n | B_n | C_n | D_n | E_6 | E_7 | E_8 | F_4 | G_2 |
| $\hat{\Gamma}$ | $A_n^{(1)}$ | $B_n^{(1)}$ | $C_n^{(1)}$ | $D_n^{(1)}$ | $E_6^{(1)}$ | $E_7^{(1)}$ | $E_8^{(1)}$ | $F_4^{(1)}$ | $G_2^{(1)}$ |
| ${}^L\Gamma$ | A_n | C_n | B_n | D_n | E_6 | E_7 | E_8 | F_4 | G_2 |
| ${}^L\hat{\Gamma}$ | $A_n^{(1)}$ | $A_{2n-1}^{(2)}$ | $D_{n+1}^{(2)}$ | $D_n^{(1)}$ | $E_6^{(1)}$ | $E_7^{(1)}$ | $E_8^{(1)}$ | $E_6^{(2)}$ | $D_4^{(3)}$ |

From this table, we see that if Γ is simply-laced, then $\hat{\Gamma} = {}^L\hat{\Gamma}$. If Γ is non simply-laced, then ${}^L\hat{\Gamma}$ is of twisted affine type.

Let ${}^L\tilde{\mathfrak{h}}$ denote the Cartan subalgebra of ${}^L\tilde{\mathfrak{g}}$. We can write

$${}^L\tilde{\mathfrak{h}} = {}^L\mathfrak{h} \oplus \mathbb{C}\check{c} \oplus \mathbb{C}\check{d}, \text{ and } {}^L\tilde{\mathfrak{h}}^* = {}^L\mathfrak{h}^* \oplus \mathbb{C}\check{\Lambda}_0 \oplus \mathbb{C}\check{\delta},$$

where ${}^L\mathfrak{h}$ is the Cartan subalgebra of ${}^L\mathfrak{g}$, \check{c} is the canonical center of ${}^L\tilde{\mathfrak{g}}$, \check{d} is the degree operator, $\check{\Lambda}_0$ is the fundamental weight of ${}^L\tilde{\mathfrak{g}}$ associated to $0 \in \hat{I}$, and $\check{\delta}$ is the element such that $\check{\delta}|_{{}^L\tilde{\mathfrak{h}}} = 0$, $\langle \check{\delta}, \check{c} \rangle = 0$ and $\langle \check{\delta}, \check{d} \rangle = 1$.

Recall that $\tilde{\mathfrak{h}}$ is the Cartan subalgebra of $\tilde{\mathfrak{g}}$. Under the duality of $\hat{\Gamma}$ and ${}^L\hat{\Gamma}$, we may identify ${}^L\tilde{\mathfrak{h}}$ with $\tilde{\mathfrak{h}}^*$ and identify ${}^L\tilde{\mathfrak{h}}^*$ with $\tilde{\mathfrak{h}}$. Under this identification, ${}^L\mathfrak{h} = \mathfrak{h}^*$ and ${}^L\mathfrak{h}^* = \mathfrak{h}$. In particular, the simple roots $\{\alpha_i | i \in \hat{I}\}$ of $\tilde{\mathfrak{g}}$ can be regarded as the simple coroots of ${}^L\tilde{\mathfrak{g}}$, and the simple coroots $\{\check{\alpha}_i | i \in \hat{I}\}$ of $\tilde{\mathfrak{g}}$ can be regarded as simple roots of ${}^L\tilde{\mathfrak{g}}$. Moreover, δ can be regarded as the canonical center \check{c} , hence

(19)

$$\check{c} = \sum_{i \in \hat{I}} a_i \alpha_i,$$

where a_i is the Kac labeling of $\hat{\Gamma}$ at i , in particular $a_0 = 1$. The coroot lattice \check{Q} (resp. coweight lattice \check{P}) are now regarded as root lattice (resp. weight lattice) of ${}^L\mathfrak{g}$.

We will still denote by $(\cdot | \cdot)$ the induced bilinear forms on ${}^L\tilde{\mathfrak{h}}$ and ${}^L\tilde{\mathfrak{h}}^*$ from the normalized bilinear form $(\cdot | \cdot)$ on $\tilde{\mathfrak{h}}$ and $\tilde{\mathfrak{h}}$. It turns out that the induced forms on ${}^L\tilde{\mathfrak{h}}$ and ${}^L\tilde{\mathfrak{h}}^*$ are still the normalized bilinear forms with respect to ${}^L\tilde{\mathfrak{g}}$.

Recall the affine Weyl group $W_{\text{aff}} = \check{Q} \rtimes W$. We can also realize W_{aff} as the Weyl group of ${}^L\tilde{\mathfrak{g}}$. For any $\tau_{\lambda} w \in W_{\text{aff}}$ and $h \in {}^L\tilde{\mathfrak{h}}^*$, following (Ka, §6.5.5) we define

(20)

$$\tau_{\lambda} w(h) = h + \langle h, \check{c} \rangle \lambda - ((h|\lambda) + \frac{(\lambda|\lambda)}{2} \langle h, \check{c} \rangle) \check{\delta}.$$

The set ${}^L\hat{\Phi}_{\text{re}}$ of real roots of ${}^L\tilde{\mathfrak{g}}$ can be described as follows (cf.(Ka, Prop.6.3 a)b))

$${}^L\hat{\Phi}_{\text{re}} = \{\check{\alpha} + kr_{\alpha}\check{\delta} | \alpha \in \Phi, k \in \mathbb{Z}\}$$

where $\check{\alpha}$ is the coroot associated to α , and $r_\alpha = \frac{2}{(\alpha|\alpha)}$, more precisely

$$r_\alpha := \begin{cases} 1 & \text{if } \Gamma \text{ is simply-laced, or } \alpha \text{ is a long root when } \Gamma \text{ is non simply-laced} \\ 2 & \text{if } \alpha \text{ is a short root when } \Gamma = B_n, C_n, F_4 \\ 3 & \text{if } \alpha \text{ is a short root when } \Gamma = G_2 \end{cases}.$$

There is a bijection $\eta : \hat{\Phi}_{\text{re}} \rightarrow {}^L\hat{\Phi}_{\text{re}}$ between $\hat{\Phi}_{\text{re}}$ and ${}^L\hat{\Phi}_{\text{re}}$, given by

$$\eta(\alpha + k\delta) = \check{\alpha} + kr_\alpha\check{\delta}.$$

Lemma 3.1. *The bijection η is W_{aff} -equivariant.*

Proof. For any $\tau_\lambda w \in W_{\text{aff}}$ and $\alpha + k\delta \in \hat{\Phi}$, from the formula (13) we have

$$\begin{aligned} \eta(\tau_\lambda w(\alpha + k\delta)) &= \eta(w(\check{\alpha}) + (k - \langle \lambda, w(\alpha) \rangle)\delta) \\ &= w(\check{\alpha}) + (k - \langle \lambda, w(\alpha) \rangle)r_\alpha\check{\delta}, \end{aligned}$$

where $w(\check{\alpha})$ is equal to the coroot of $w(\alpha)$. On the other hand,

$$\begin{aligned} \tau_\lambda w(\eta(\alpha + k\delta)) &= \tau_\lambda w(\check{\alpha} + kr_\alpha\check{\delta}) \\ &= w(\check{\alpha}) + (kr_\alpha - \langle \lambda, w(\check{\alpha}) \rangle)\check{\delta} \\ &= w(\check{\alpha}) + (kr_\alpha - \langle \lambda, w(\check{\alpha}) \rangle)r_\alpha\check{\delta}, \end{aligned}$$

where the second equality follows from (20), and the third equality holds since

$$\langle \lambda, w(\check{\alpha}) \rangle = \frac{2\langle \lambda, w(\check{\alpha}) \rangle}{(w(\alpha)|w(\alpha))} = \frac{2\langle \lambda, w(\check{\alpha}) \rangle}{(\alpha|\alpha)} = \langle \lambda, w(\check{\alpha}) \rangle r_\alpha.$$

From the above two computations, we see that η is W_{aff} -equivariant. \square

Under the map η , the image $\eta(\hat{\Phi}_{\text{re}}^\pm)$ is the set of all positive (resp. negative) roots in ${}^L\hat{\Phi}_{\text{re}}$. Recall the set $\{\check{\omega}_\kappa \mid \kappa \in \hat{M}\}$ in Lemma 2.14. The following lemma follows from the discussion in (Ka, §12.4). For the convenience of the reader, we include an argument here.

Lemma 3.2. *For any $\lambda \in \check{P}^+$, $\check{\Lambda}_0 + \lambda$ is a dominant weight of ${}^L\check{\mathfrak{g}}$ of level one if and only if $\lambda \in \{\check{\omega}_\kappa \mid \kappa \in \hat{M}\}$.*

Proof. We first observe that λ is dominant if and only if

$$\langle \check{\Lambda}_0 + \lambda, \alpha_i \rangle = \langle \lambda, \alpha_i \rangle \geq 0, \quad \text{for any } i \in I.$$

By the formulae (19) and (14), $\alpha_0 = \check{c} - \theta$. Hence, $\check{\Lambda}_0 + \lambda$ is dominant if and only

$$\langle \check{\Lambda}_0 + \lambda, \alpha_0 \rangle = \langle \check{\Lambda}_0 + \lambda, \check{c} - \theta \rangle = 1 - \langle \lambda, \theta \rangle \geq 0.$$

By Lemma 2.13, λ is equal to $\check{\omega}_\kappa$ for $\kappa \in \hat{M}$. This concludes the proof. \square

For each $\kappa \in \hat{M}$, set $\check{\Lambda}_\kappa := \check{\Lambda}_0 + \check{\omega}_\kappa$.

Lemma 3.3. *The stabilizer group of W_{aff} at the dominant weight $\check{\Lambda}_\kappa$ is equal to W_κ .*

Proof. It is known that the stabilizer W''_κ of W_{aff} at $\check{\Lambda}_\kappa$ is a parabolic subgroup of W_{aff} . It is enough to determine the Coxeter generators of W''_κ . We first examine $s_0(\check{\Lambda}_0)$,

$$s_0(\check{\Lambda}_0) = \tau_{\check{\theta}} s_{\check{\theta}}(\check{\Lambda}_0) = \tau_{\check{\theta}}(\check{\Lambda}_0) = \check{\Lambda}_0 + \check{\theta} - \frac{(\check{\theta}|\check{\theta})}{2}\check{\delta} = \check{\Lambda}_0 + \check{\theta} - \check{\delta}.$$

For any $\kappa \in M$,

$$\begin{aligned} s_0(\check{\Lambda}_\kappa) &= \tau_{\check{\theta}} s_{\check{\theta}}(\check{\Lambda}_0 + \check{\omega}_\kappa) = \tau_{\check{\theta}}(\check{\Lambda}_0 + \check{\omega}_\kappa - \check{\theta}) \\ &= \check{\Lambda}_0 + \check{\omega}_\kappa - \check{\theta} + \langle \check{\Lambda}_0 + \check{\omega}_\kappa - \check{\theta}, \check{c} \rangle \check{\theta} - ((\check{\theta}|\check{\Lambda}_0 + \check{\omega}_\kappa - \check{\theta}) + \frac{(\check{\theta}|\check{\theta})}{2} \langle \check{\Lambda}_0 + \check{\omega}_\kappa, \check{c} \rangle) \check{\delta} \\ &= \check{\Lambda}_0 + \check{\omega}_\kappa - ((\check{\theta}|\check{\omega}_\kappa - \check{\theta}) + \frac{(\check{\theta}|\check{\theta})}{2}) \check{\delta} = \check{\Lambda}_\kappa, \end{aligned}$$

where the fourth equality holds since $(\check{\theta}|\check{\theta}) = 2$ and $(\check{\theta}|\check{\omega}_\kappa - \check{\theta}) = -1$.

Now it is easy to see that the Coxeter generators of W''_κ are given by $\{s_i \mid i \in I_\kappa\}$, where $I_\kappa = \hat{I} \setminus \{\kappa\}$. By Proposition 2.15, we can conclude that $W''_\kappa = W_\kappa$. □

3.2. Level one affine Demazure Modules. We first make a digression to prove a general lemma in the setting of general symmetrizable Kac-Moody algebras. Let Λ be an integral dominant weight of a symmetrizable Kac-Moody algebra \mathcal{G} with a fixed Borel subalgebra \mathcal{B} . Let \mathcal{W} be the Weyl group of \mathcal{G} . Let V_Λ denote the irreducible integrable representation of \mathcal{G} of highest weight Λ . Let \mathcal{W}_Λ denote the stabilizer group of \mathcal{W} at Λ . Then \mathcal{W}_Λ is a parabolic subgroup of \mathcal{W} . Let \mathcal{W}^Λ denote the set of minimal coset representatives in $\mathcal{W}/\mathcal{W}_\Lambda$.

Proposition 3.4. *For any $y, w \in \mathcal{W}^\Lambda$, we have*

$$w < y \iff v_{w(\Lambda)} \in U(\mathcal{B})v_{y(\Lambda)}$$

Proof. Let \mathcal{B}^- denote the negative Borel subalgebra in \mathcal{G} . Then we have

$$(21) \quad w < y \iff v_{y(\Lambda)} \in U(\mathcal{B}^-)v_{w(\Lambda)}.$$

This fact is stated in (Ku, ex.7.1.E.4), and this is a parabolic version of (Ku, Prop. 7.1.20). The proof is almost identical to that in (Ku, Prop. 7.1.20).

In the next step, we show that

$$v_{y(\Lambda)} \in U(\mathcal{B}^-)v_{w(\Lambda)} \iff v_{w(\Lambda)} \in U(\mathcal{B})v_{y(\Lambda)}.$$

Let (\cdot, \cdot) be the contravariant form on \mathcal{H}_Λ (cf. (Ku, Prop. 2.3.2)), which satisfies that, for any $v_1, v_2 \in V_\Lambda$, and $x \in \mathcal{G}$.

$$(xv_1, v_2) = (v_1, \sigma(x)v_2),$$

where $\sigma : \mathcal{G} \rightarrow \mathcal{G}$ is the Cartan involution on \mathcal{G} . The involution σ induces an anti-automorphism on the universal enveloping algebra $U(\mathcal{G})$ of \mathcal{G} . Furthermore the contravariant form (\cdot, \cdot) has the following properties:

- (1) (\cdot, \cdot) is non-degenerate on each weight space $V_\Lambda(\mu)$, where μ is a weight of \mathcal{G} .
- (2) $(v_1, v_2) = 0$ for any two weight vectors v_1, v_2 of distinct weights.

Assume that $v_{y(\Lambda)} \in U(\mathcal{B}^-)v_{w(\Lambda)}$. We may write $v_{y(\Lambda)} = Pv_{w(\Lambda)}$, where P is an element in the enveloping algebra $U(\mathcal{N}^-)$ of the nilpotent radical \mathcal{N}^- of \mathcal{B}^- . Notice that the weight space $V_\Lambda(y(\Lambda))$ is 1-dimensional. It follows that we may assume $P \in U(\mathcal{N}^-)$ is a monomial in negative root vectors in \mathcal{N}^- . By the non-degeneracy of (\cdot, \cdot) on $V_\Lambda(y(\Lambda))$, we have

$$0 \neq (v_{y(\Lambda)}, v_{y(\Lambda)}) = (Pv_{w(\Lambda)}, v_{y(\Lambda)}) = (v_{w(\Lambda)}, \sigma(P)v_{y(\Lambda)}).$$

Note that $\sigma(P) \in U(\mathcal{N})$ is a monomial in positive root vectors in \mathcal{N} , where \mathcal{N} is the unipotent radical of \mathcal{B} . By the second property of the contravariant form mentioned above, we must have $\sigma(P)v_{y(\Lambda)} \in V_\Lambda(w(\Lambda))$. By the one-dimensionality of $V_\Lambda(w(\Lambda))$, there exists a nonzero constant c such that

$$v_{w(\Lambda)} = c\sigma(P)v_{y(\Lambda)} \in U(\mathcal{N})v_{y(\Lambda)}.$$

By similar argument, we can show that if $v_{w(\Lambda)} \in U(\mathcal{B})v_{y(\Lambda)}$, then $v_{y(\Lambda)} \in U(\mathcal{B}^-)v_{w(\Lambda)}$. Thus the lemma is proven. \square

Remark 3.5. *The equivalence (21) can be proved by the induction on the length of elements of Weyl group. The proof of (Ku, Prop. 7.1.20) in the regular case essentially incorporates the original proof for finite Weyl groups by Bernstein-Gelfand-Gelfand (BGG). We don't know how to prove Proposition 3.4 by induction directly. It is interesting to use the contravariant form to reduce the lemma to the equivalence (21).*

We now return to the Kac-Moody algebra ${}^L\check{\mathfrak{g}}$. Let \mathcal{H}_κ denote the irreducible integrable representation of ${}^L\check{\mathfrak{g}}$ of highest weight $\check{\Lambda}_\kappa$. Then \mathcal{H}_κ is an integrable representation of level one. Fix a highest weight vector $v_{\check{\Lambda}_\kappa} \in \mathcal{H}_\kappa$. For any $\lambda \in \check{P}$ if $\lambda \in \check{Q}_\kappa := \check{Q} - \check{\omega}_\kappa$, then we set

$$\varpi(\lambda) := \tau_{-\lambda-\check{\omega}_\kappa}(\check{\Lambda}_\kappa),$$

where $\tau_{-\lambda-\check{\omega}_\kappa} \in W_{\text{aff}}$. Let $v_{\varpi(\lambda)} \in \mathcal{H}_\kappa$ be an extremal vector of weight $\varpi(\lambda)$. By this convention, $\varpi(-\check{\omega}_\kappa) = \check{\Lambda}_\kappa$. By a simple computation from formula (20), we have the following formula

$$(22) \quad \varpi(\lambda) = \check{\Lambda}_0 - \lambda - \frac{(\lambda|\lambda) - (\check{\omega}_\kappa|\check{\omega}_\kappa)}{2}\check{\delta}, \quad \text{for } \lambda \in \check{Q}_\kappa.$$

Definition 3.6. We now define the affine Demazure module $\hat{\mathcal{D}}_\lambda$ for each $\lambda \in \check{P}$ as follows,

$$\hat{\mathcal{D}}_\lambda := U({}^L\check{\mathfrak{b}})v_{\varpi(\lambda)} \subset \mathcal{H}_\kappa, \quad \text{if } \lambda \in \check{Q}_\kappa,$$

where $U({}^L\check{\mathfrak{b}})$ is the universal enveloping algebra of the Borel subalgebra ${}^L\check{\mathfrak{b}}$ of ${}^L\check{\mathfrak{g}}$.

Recall that there is a bijection $\check{Q}_\kappa \simeq W_{\text{aff}}/W_\kappa$ where $\check{Q}_\kappa := \check{Q} - \check{\omega}_\kappa$. For each $\lambda \in \check{Q}_\kappa$, let $\bar{\tau}_{-\lambda-\check{\omega}_\kappa}$ denote the associated minimal representative in the coset $\tau_{-\lambda-\check{\omega}_\kappa}W_\kappa$.

Proposition 3.7. *For any $\lambda, \mu \in \check{P}$, then $\mu <_I \lambda$ if and only if $v_{\varpi(\mu)} \in \hat{\mathcal{D}}_\lambda$.*

Proof. First of all, $\mu <_I \lambda$ if and only if $\bar{\tau}_{-\lambda-\check{\omega}_\kappa} < \bar{\tau}_{-\mu-\check{\omega}_\kappa}$. By Lemma 3.3, we have

$$\varpi(\lambda) = \bar{\tau}_{-\lambda-\check{\omega}_\kappa}(\check{\Lambda}_\kappa), \quad \varpi(\mu) = \bar{\tau}_{-\mu-\check{\omega}_\kappa}(\check{\Lambda}_\kappa).$$

Hence $\hat{\mathcal{D}}_\lambda = U({}^L\check{\mathfrak{b}}) \cdot v_{\varpi(\lambda)}$. Lastly, in view of Proposition 3.4 we conclude that $\mu <_I \lambda$ if and only if $v_{\varpi(\mu)} \in \hat{\mathcal{D}}_\lambda$. \square

Let $P(\mathcal{H}_\kappa)$ denote the weight system of the integrable representation \mathcal{H}_κ . A weight $\hat{\mu} \in P(\mathcal{H}_\kappa)$ is called maximal if $\hat{\mu} + \check{\delta} \notin P(\mathcal{H}_\kappa)$. Let $\text{Max}(\mathcal{H}_\kappa)$ denote the set of all maximal weights in $P(\mathcal{H}_\kappa)$. The weight system $P(\mathcal{H}_\kappa)$ can be completely described by the following lemma (cf. (Ka, §12.6)).

Lemma 3.8. *For any $\kappa \in \hat{M}$, we have*

- (1) $\text{Max}(\mathcal{H}_\kappa) = \{\varpi(\lambda) \mid \lambda \in \check{Q}_\kappa\}$;
- (2) $P(\mathcal{H}_\kappa) = \bigcup_{\lambda \in \check{Q}_\kappa} \{\varpi(\lambda) - n\check{\delta} \mid n \in \mathbb{Z}^+\}$ is a disjoint union.

Any weight in $P(\mathcal{H}_\kappa)$ is of the form $\check{\Lambda}_0 - \lambda + m\check{\delta}$, for some integer m . We define a twisted version of projection

$$\mathfrak{p} : P(\mathcal{H}_\kappa) \rightarrow \check{P}, \text{ given by } \check{\Lambda}_0 - \lambda + m\check{\delta} \mapsto \lambda.$$

Let $P(\hat{\mathcal{D}}_\lambda)$ denote the weight system of the affine Demazure module $\hat{\mathcal{D}}_\lambda$.

Theorem 3.9. *The map \mathfrak{p} maps $P(\hat{\mathcal{D}}_\lambda)$ onto $\Psi(\lambda)$, and \mathfrak{p} admits a canonical section $\lambda \mapsto \varpi(\lambda)$.*

Proof. Set

$$\mathfrak{H} := \bigoplus_{n>0} ({}^L\check{\mathfrak{g}})_{\pm n\check{\delta}} \oplus {}^L\mathfrak{h} \oplus \mathbb{C}\check{c}, \text{ and } \mathfrak{H}^\pm := \bigoplus_{n>0} ({}^L\check{\mathfrak{g}})_{\pm n\check{\delta}}.$$

Then \mathfrak{H} is an Heisenberg algebra with the center \check{c} . By the Frenkel-Kac construction (for untwisted affine types, see (FK); for $A_{2n-1}^{(2)}, D_{n+1}^{(2)}, E_6^{(2)}$, see (Fr, Theorem I.2.25) (FLM, §7); for $D_4^{(3)}$, see (BT, §B.8)), the representation \mathcal{H}_κ can be realized as

$$(23) \quad \mathcal{H}_\kappa \simeq S(\mathfrak{H}^-) \otimes \mathbb{C}[\check{Q}_\kappa],$$

where $S(\mathfrak{H}^-)$ is the symmetric algebra of \mathfrak{H}^- , and $\mathbb{C}[\check{Q}_\kappa]$ consists of linear combinations of e^λ , $\lambda \in \check{Q}_\kappa$. Moreover

$$\mathcal{H}_\kappa(\varpi(\lambda) - n\check{\delta}) \simeq S(\mathfrak{H}^-)_{-n\check{\delta}} \otimes e^\lambda,$$

where e^λ is of weight $\varpi(\lambda)$, and

$$\bigoplus_{n \geq 0} \mathcal{H}_\kappa(\varpi(\lambda) - n\check{\delta}) \simeq S(\mathfrak{H}^-) \otimes e^\lambda$$

is a free $U(\mathfrak{H}^-)$ -module of rank 1.

By Lemma 3.8, any weight vector in $\hat{\mathcal{D}}_\lambda$ is of weight $\varpi(\mu) - m\check{\delta}$ for some integer m and $\mu \in \check{P}$. To show that \mathfrak{p} maps $P(\hat{\mathcal{D}}_\lambda)$ onto $\Psi(\lambda)$, it suffices to show that for some nonzero weight vector $x_{\varpi(\mu) - m\check{\delta}}$ in $\hat{\mathcal{D}}_\lambda$ of weight $\varpi(\mu) - m\check{\delta}$, the maximal vector $v_{\varpi(\mu)}$ is also an element in $\hat{\mathcal{D}}_\lambda$.

Let (\cdot, \cdot) be the contravariant form on \mathcal{H}_κ , and let σ be the Cartan involution on ${}^L\check{\mathfrak{g}}$. Clearly, there exists a weight vector $x_{\varpi(\mu) - m\check{\delta}}$ of weight $\varpi(\mu) - m\check{\delta}$, such that

$$(x_{\varpi(\mu) - m\check{\delta}}, x_{\varpi(\mu) - m\check{\delta}}) \neq 0,$$

since the contravariant form (\cdot, \cdot) on the weight space $\mathcal{H}_\kappa(\varpi(\mu) - m\check{\delta})$ is nondegenerate. By the construction in (23), $x_{\varpi(\mu) - m\check{\delta}}$ can be written as $P \cdot v_{\varpi(\mu)}$, where P is an element of weight $-m\check{\delta}$ in the enveloping algebra $U(\mathfrak{H}^-)$ of \mathfrak{H}^- . Then


$$0 \neq (x_{\varpi(\mu) - m\check{\delta}}, x_{\varpi(\mu) - m\check{\delta}}) = (P \cdot v_{\varpi(\mu)}, x_{\varpi(\mu) - m\check{\delta}}) = (v_{\varpi(\mu)}, \sigma(P)x_{\varpi(\mu) - m\check{\delta}}).$$

By the one-dimensionality of the weight space $\mathcal{H}_k(\varpi(\mu))$, $\sigma(P)x_{\varpi(\mu)-m\check{\delta}} = cv_{\varpi(\mu)}$ for some nonzero constant $c \in \mathbb{C}$, since $\sigma(P) \in U(\mathfrak{H}^+)_{m\check{\delta}}$. Here we use the fact that σ maps \mathfrak{H}^- to \mathfrak{H}^+ . When ${}^L\tilde{\mathfrak{g}}$ is of untwisted affine types, this fact follows from (Ka, §7.6). This fact holds for twisted affine types as well, see (Ka, §8.3).

Since $\mathfrak{H}^+ \subset {}^L\tilde{\mathfrak{b}}$, it follows that $v_{\varpi(\mu)} \in \hat{\mathcal{D}}_\lambda$. Then by Proposition 3.7, we have $\mu <_I \lambda$. In other words, $\mu \in \Psi(\lambda)$. This concludes the proof. \square

Remark 3.10. *In the proof of Theorem 3.9, we crucially use the Frenkel-Kac construction of basic representations for ${}^L\tilde{\mathfrak{g}}$. Note that this construction only works for affine Kac-Moody algebras of type $X_n^{(r)}$, where $r = 1, 2, 3$ and X_n is of type A, D, E. From Table (18), we see that ${}^L\tilde{\mathfrak{g}}$ exhausts all cases except $A_{2n}^{(2)}$.*

Example 3.11. By Lemma 3.8, it is clear that $\mathfrak{p} : \mathbb{P}(\mathcal{H}_\kappa) \rightarrow \check{P}$ is not one-to-one. In fact $\mathfrak{p} : \mathbb{P}(\hat{\mathcal{D}}_\lambda) \rightarrow \Psi(\lambda)$ is not one-to-one as well. For example when $\mathfrak{g} = sl_2$ and $\kappa = 0$, consider the affine Demazure module $\hat{\mathcal{D}}_{-2\check{\alpha}}$, where α is the simple root of \mathfrak{g} . One can check that $0 \neq (e_{\check{\alpha}})^2 v_{\varpi(-2\check{\alpha})}$ has weight $\check{\Lambda}_0 - 2\check{\delta}$. The following example describes all weights appearing in $\hat{\mathcal{D}}_{-2\check{\alpha}}$.



Let $e_{\check{\alpha}+mr_\alpha\check{\delta}}$ be a root vector in ${}^L\tilde{\mathfrak{g}}$ corresponding to the root $\check{\alpha} + mr_\alpha\check{\delta} \in {}^L\hat{\Phi}^+$. Via the bijection η given in Section 3.1, by Lemma 3.1 the coroot of $\check{\alpha} + mr_\alpha\check{\delta}$ is $\alpha + m\delta \in \hat{\Phi}^+$.

The following lemma gives a representation theoretic interpretation for Lemma 2.2.

Lemma 3.12. *For any $\alpha \in \Phi^+$,*

(1) *if $\langle \lambda, \alpha \rangle > 0$, then for any $0 < k \leq \langle \lambda, \alpha \rangle$,*

$$(24) \quad (e_{\check{\alpha}})^k \cdot v_{\varpi(\lambda)} \neq 0, \text{ and } \mathfrak{p}(\text{wt}((e_{\check{\alpha}})^k \cdot v_{\varpi(\lambda)})) = \lambda - k\check{\alpha};$$

(2) *if $\langle \lambda, \alpha \rangle < -1$, then for any $0 < k < -\langle \lambda, \alpha \rangle$,*

$$(25) \quad (e_{-\check{\alpha}+r_\alpha\check{\delta}})^k \cdot v_{\varpi(\lambda)} \neq 0, \text{ and } \mathfrak{p}(\text{wt}(e_{-\check{\alpha}+r_\alpha\check{\delta}})^k \cdot v_{\varpi(\lambda)}) = \lambda + k\check{\alpha},$$

where $\text{wt}(\cdot)$ denote the weight of a weight vector.

Proof. With respect to any sl_2 -triple $\{e_{\check{\alpha}+kr_\alpha\check{\delta}}, e_{-\check{\alpha}-kr_\alpha\check{\delta}}, \alpha+k\delta\}$ associated to $\check{\alpha}+kr_\alpha\check{\delta} \in \hat{\Phi}^+$, the extremal vector $v_{\varpi(\lambda)}$ is either a highest weight vector or lowest weight vector. In part (1), by (22)

$$\langle \varpi(\lambda), \alpha \rangle = -\langle \lambda, \alpha \rangle < 0,$$

the vector $v_{\varpi(\lambda)}$ is a weight vector of lowest weight $-\langle \lambda, \alpha \rangle$ with respect to the sl_2 -triple associated to $\check{\alpha}$. Hence the statement (24) holds.

In part (2), since

$$\langle \varpi(\lambda), -\alpha + \delta \rangle = 1 + \langle \lambda, \alpha \rangle < 0,$$

the vector $v_{\varpi(\lambda)}$ is again a weight vector of lowest weight $1 + \langle \lambda, \alpha \rangle$ with respect to the sl_2 -triple associated to the positive root $-\check{\alpha} + r_\alpha \check{\delta}$. Hence the statement (25) holds as well. \square

Remark 3.13. For any $\lambda \in \check{Q}_\kappa$, let γ denote the unique dominant coweight that is translated by the Weyl group W from $-\lambda$. Set $V_\gamma := U({}^L\mathfrak{g}) \cdot v_{\varpi(\lambda)} \subset \mathcal{H}_\kappa$. It is well-known that V_γ is a finite dimensional irreducible representation of highest weight γ . The vector $v_{\varpi(\lambda)} \in V_\gamma$ is an extremal vector of weight $-\lambda$. Let ${}^L\mathfrak{b}$ denote the Borel subalgebra of ${}^L\mathfrak{g}$ obtained as ${}^L\mathfrak{b} := {}^L\check{\mathfrak{b}} \cap {}^L\mathfrak{g}$. Then $D_{-\lambda} := U({}^L\mathfrak{b})v_{\varpi(\lambda)}$ is a Demazure module in V_γ in the usual sense. It is proved in (Io2) that the multiplicity of the weight space $D_{-\lambda}(-\mu)$ is equal to the number of top components of the intersection $N(\mathcal{X}) \cdot L_\mu \cap I \cdot L_\lambda$. It is a natural question to ask if there is similar phenomenon for the affine Demazure module $\hat{\mathcal{D}}_\lambda$.

4. FURTHER STUDY OF THE SET $\Psi(\lambda)$

4.1. The partial order $<_I$ on a chamber of the coweight lattice. Let \mathfrak{C} be the dominant chamber of the coweight lattice \check{P} determined by $T \subset B$, i.e.

$$\mathfrak{C} = \{\lambda \in \check{P} \mid \langle \lambda, \alpha \rangle \geq 0, \text{ for any } \alpha \in \Phi^+\}.$$

Set $\mathfrak{C}_w := w(\mathfrak{C})$ for any $w \in W$. Then \check{P} is the union of all chambers \mathfrak{C}_w indexed by $w \in W$.

Theorem 4.1. For any coweights $\lambda, \mu \in \mathfrak{C}_w$, $\mu <_I \lambda$ if and only if $\lambda - \mu$ is a positive sum of coroots in $w(\check{\Phi}^+)$.

Proof. We first assume that $\mu <_I \lambda$, in other words $L_\mu \subset \bar{\mathbf{X}}_\lambda$. Set $\lambda^+ := w^{-1}(\lambda)$ and $\mu^+ := w^{-1}(\mu)$. Then $\lambda^+, \mu^+ \in \mathfrak{C}$. It follows that $L_{\mu^+} \subset \overline{\mathbf{Gr}}_{\lambda^+}$, since $\mathbf{Gr}_{\lambda^+} = G(\mathcal{O}) \cdot L_{\lambda^+}$. Hence $\lambda^+ - \mu^+$ is a positive sum of coroots in $\check{\Phi}^+$. Equivalently, $\lambda - \mu$ is a positive sum of coroots in $w(\check{\Phi}^+)$.

Now we prove the converse. Assume that $\lambda - \mu$ is a positive sum of coroots in $w(\check{\Phi}^+)$. Equivalently, $\lambda^+ - \mu^+$ is a positive sum of coroots in $\check{\Phi}^+$. By a result due to Stembridge and Steinberg (cf. (Stg, Corollary 2.7)), there exists a sequence of dominant coweights

$$\mu^+ = \lambda_0^+, \lambda_1^+, \dots, \lambda_{k-1}^+, \lambda_k^+ = \lambda^+,$$

such that for each $1 \leq i \leq k$, $\lambda_i^+ = \lambda_{i-1}^+ + \check{\beta}_i$ where $\check{\beta}_i \in \check{\Phi}^+$. Set $\lambda_i := w(\lambda_i^+)$ for each i . Then $\lambda_i = \lambda_{i-1} + w(\check{\beta}_i)$. Since

$$\langle \lambda_i, w(\beta_i) \rangle = \langle \lambda_{i-1} + w(\check{\beta}_i), w(\beta_i) \rangle = \langle \lambda_{i-1}^+ + \check{\beta}_i, \beta_i \rangle = \langle \lambda_{i-1}^+, \beta_i \rangle + 2 > 0,$$

by Lemma 2.2, $\lambda_{i-1} = \lambda_i - w(\check{\beta}_i) \in \Psi(\lambda_i)$. Hence $\lambda_{i-1} <_I \lambda_i$. Inductively, we have $\mu <_I \lambda$. \square

This theorem completely describes the partial order $<_I$ on any fixed chamber. Moreover from this theorem, we immediately get the following corollary.

Corollary 4.2. *For any two coweights λ, μ in the same chamber, $\mu <_I \lambda$ if and only if $w(\mu) <_I w(\lambda)$.*

4.2. R-operators and cover relations. For each $\alpha \in \Phi^+$, we define the operator R_α on \check{P} ,

$$(26) \quad R_\alpha(\lambda) = \begin{cases} s_\alpha(\lambda) := \lambda - \langle \lambda, \alpha \rangle \check{\alpha}, & \text{if } \langle \lambda, \alpha \rangle \geq 0 \\ s_\alpha(\lambda) - \check{\alpha} =: \lambda - \langle \lambda, \alpha \rangle \check{\alpha} - \alpha, & \text{if } \langle \lambda, \alpha \rangle < 0 \end{cases}.$$

By Lemma 2.2, for any $\lambda \in \check{P}$, we always have $R_\alpha(\lambda) \in \Psi(\lambda)$.

The following theorem shows that the set $\Psi(\lambda)$ can be obtained by repeating applying R-operators starting at λ .

Theorem 4.3. *For any coweight $\lambda \in \check{P}$, we have*

$$\Psi(\lambda) = \{R_{\beta_1} R_{\beta_2} \cdots R_{\beta_k}(\lambda) \mid k \in \mathbb{N}, \beta_1, \beta_2, \dots, \beta_k \in \Phi^+\}.$$

Proof. By Lemma 2.2, for any $\lambda \in \check{P}$, we always have $R_\alpha(\lambda) <_I \lambda$. Therefore for any sequence of positive roots β_1, \dots, β_k , we have $R_{\beta_1} R_{\beta_2} \cdots R_{\beta_k}(\lambda) \in \Psi(\lambda)$.

To show the other inclusion, we first prove the following general fact: for any $\mu, \mu' \in \check{P}$, if $\mu' \in S(\mu, \alpha)$ for some positive root $\alpha \in \Phi^+$ (recall $S(\mu, \alpha)$ defined in (10)), then μ' can be written as $(R_\alpha)^k(\mu)$ for some integer k . We first assume that $\langle \mu, \alpha \rangle \geq 0$. One can check that for any integer $m \geq 0$,

$$(27) \quad \mu - m\check{\alpha} = \begin{cases} (R_\alpha)^{2m}(\mu), & \text{if } 0 \leq m \leq \frac{1}{2}\langle \mu, \alpha \rangle \\ (R_\alpha)^{2(\langle \mu, \alpha \rangle - m) + 1}(\mu), & \text{if } \frac{1}{2}\langle \mu, \alpha \rangle < m \leq \langle \mu, \alpha \rangle \end{cases}.$$

Now we assume that $\langle \mu, \alpha \rangle < 0$, one can check similarly

$$(28) \quad \mu + m\check{\alpha} = \begin{cases} (R_\alpha)^{2m}(\mu), & \text{if } 0 \leq m < -\frac{1}{2}\langle \mu, \alpha \rangle \\ (R_\alpha)^{2(-\langle \mu, \alpha \rangle - m) - 1}(\mu), & \text{if } -\frac{1}{2}\langle \mu, \alpha \rangle \leq m \leq -\langle \mu, \alpha \rangle - 1 \end{cases}.$$

From the computations (27) (28), we see that this theorem follows from Theorem 2.6. \square

For $\alpha \in \Phi^+$ and $k \in \mathbb{Z}$, let $H_{\alpha, k}$ denote the hyperplane $\{\lambda \in \check{P} \mid \langle \lambda, \alpha \rangle = k\}$.

Lemma 4.4. *Assume that μ, λ are in the same chamber \mathfrak{C}_w and $\mu, \lambda \notin H_{\alpha, -1}$ for some positive root $\alpha \in \Phi^+$, if $\mu <_I \lambda$, then $R_\alpha(\mu) <_I R_\alpha(\lambda)$.*

Proof. Since α is either an element in $w(\Phi^+)$ or an element in $w(\Phi^-)$, it follows that either $\langle \lambda, \alpha \rangle \geq 0$ and $\langle \mu, \alpha \rangle \geq 0$, or $\langle \lambda, \alpha \rangle \leq 0$ and $\langle \mu, \alpha \rangle \leq 0$. By the assumption $\mu, \lambda \notin H_{\alpha, -1}$, it follows that either $\langle \lambda, \alpha \rangle \geq 0$ and $\langle \mu, \alpha \rangle \geq 0$, or $\langle \lambda, \alpha \rangle < 0$ and $\langle \mu, \alpha \rangle < 0$.

In the first case $R_\alpha(\mu) <_I R_\alpha(\lambda)$, which follows from Corollary 4.2. In the second case,

$$R_\alpha(\lambda) = s_\alpha(\lambda) - \check{\alpha}, \text{ and } R_\alpha(\mu) = s_\alpha(\mu) - \check{\alpha}.$$

Hence by Theorem 4.1, $R_\alpha(\lambda) - R_\alpha(\mu)$ is a positive sum of coroots in $s_\alpha w(\check{\Phi}^+)$. Then by Theorem 4.1 again, $R_\alpha(\mu) <_I R_\alpha(\lambda)$. This concludes the proof. \square

In the following, we introduce the notion of α -regularity n -regularity for a coweight. This notion will allow us to cross the wall $H_\alpha := H_{\alpha,0}$.

Definition 4.5. (1) We say that a coweight λ is α -regular, if $\langle \lambda, \alpha \rangle > 0$, or $\langle \lambda, \alpha \rangle < 0$ and $\lambda + \check{\alpha} \in \mathfrak{C}_{w^\lambda}$, where w^λ is the minimal element in W such that $\lambda \in \mathfrak{C}_{w^\lambda}$.
(2) We say that λ is n -regular for a nonnegative integer n , if $\langle \lambda, w^\lambda(\alpha_i) \rangle \geq n$ for any simple root α_i .

Set

$$r = \begin{cases} 1, & \text{if } \Gamma \text{ is simply-laced;} \\ 2, & \text{if } \Gamma \text{ is non simply-laced but } \Gamma \neq G_2; \\ 3, & \text{if } \Gamma = G_2 \end{cases} .$$

Lemma 4.6. *If λ is r -regular, then λ is α -regular for any positive root α .*

Proof. By the assumption of r -regularity, it is easy to see that $\langle \lambda, \alpha \rangle \neq 0$ for any root α , since α is always a positive (or a negative) summation of $w^\lambda(\alpha_i)$. We may only consider the case when $\langle \lambda, \alpha \rangle < 0$. Note that we always have $|\langle \check{\alpha}, w^\lambda(\alpha_i) \rangle| \leq r$ (cf. (Hu1, §9.4)). It follows that

$$\langle \lambda + \check{\alpha}, w^\lambda(\alpha_i) \rangle \geq 0, \text{ for each simple root } \alpha_i.$$

Equivalently, $\lambda + \check{\alpha} \in \mathfrak{C}_{w^\lambda}$. It finishes the proof of the lemma. \square

Lemma 4.7. *Let α be a positive root in Φ^+ . Let λ be an α -regular coweight. Then $\lambda \notin H_{\alpha,-1}$ and $R_\alpha(\lambda) \in \mathfrak{C}_{s_\alpha w}$, where w is the minimal element in W such that $\lambda \in \mathfrak{C}_w$.*

Proof. First of all we show that $\lambda \notin H_{\alpha,-1}$. Assume that $\langle \lambda, \alpha \rangle \leq -1$. Then $w^{-1}(\alpha) \in \Phi^-$, since $\lambda \in \mathfrak{C}_w$. Write $-\alpha = \sum c_i w(\alpha_i)$ as a positive linear combination of the w -translations $w(\alpha_i)$ of simple roots. Then

$$\langle \lambda + \check{\alpha}, \alpha \rangle = - \sum c_i \langle \lambda + \check{\alpha}, w(\alpha_i) \rangle \leq 0.$$

By the α -regularity of λ , it follows that $\langle \lambda, \alpha \rangle \leq -2$. It follows that $\lambda \notin H_{\alpha,-1}$.

We now show that $R_\alpha(\lambda) \in \mathfrak{C}_{s_\alpha w}$. When $\langle \lambda, \alpha \rangle > 0$, it is obvious. Set $\beta := -w^{-1}(\alpha) \in \Phi^+$. Then $\check{\beta} = -w^{-1}(\check{\alpha})$. For any simple root α_i ,

$$\langle \check{\beta}, \alpha_i \rangle \leq 2$$

To show that $R_\alpha(\lambda) \in \mathfrak{C}_{s_\alpha w}$, it suffices to check that, for any simple root α_i , $\langle R_\alpha(\lambda), s_\alpha w(\alpha_i) \rangle \geq 0$, which follows since

$$\begin{aligned} \langle R_\alpha(\lambda), s_\alpha w(\alpha_i) \rangle &= \langle s_\alpha(\lambda) - \check{\alpha}, s_\alpha w(\alpha_i) \rangle \\ &= \langle \lambda + \check{\alpha}, w(\alpha_i) \rangle \geq 0. \end{aligned}$$

\square

The following result immediately follows from Lemma 4.4 and Lemma 4.7.

Proposition 4.8. *Let α be a positive root. For any α -regular coweights μ, λ in \mathfrak{C}_w , if $\mu <_I \lambda$, then $R_\alpha(\mu), R_\alpha(\lambda) \in \mathfrak{C}_{s_\alpha w}$ and $R_\alpha(\mu) <_I R_\alpha(\lambda)$.*

Following (BGG), we write $y \xrightarrow{\alpha} w$ for any $y, w \in W$, if $y = s_\alpha w$ and $\ell(w) = \ell(y) + 1$ for some reflection s_α . Similarly, we write $\mu \xrightarrow{\alpha} \lambda$ for any $\mu, \lambda \in \check{P}$, if $\mu = R_\alpha(\lambda)$ and $\dim \mathbf{X}_\lambda = \dim \mathbf{X}_\mu + 1$ for some positive root α . As shown in Theorem 4.3, the R -operators generate the closure relations for $<_I$. So we see that the covering relations of $<_I$ on \check{P} must all be of the form $\mu \xrightarrow{\alpha} \lambda$.

Recall that $\dim \mathbf{X}_\lambda = \ell(\tau_{-\lambda} w^\lambda)$ where ℓ is the length function defined in (12), and w^λ is the minimal element in W such that $\lambda \in \mathfrak{C}_{w^\lambda}$. Set $\lambda^+ := (w^\lambda)^{-1}(\lambda)$; by definition λ^+ is the dominant translate of λ . The following formula is a consequence of (12),

$$(29) \quad \dim \mathbf{X}_\lambda = 2\langle \lambda^+, \rho \rangle - \ell(w^\lambda).$$

From this formula, we immediately see that $\mu \xrightarrow{\alpha} \lambda$ if and only if $\mu = R_\alpha(\lambda)$ and

$$(30) \quad \ell(w^\lambda) - \ell(w^\mu) = 2\langle \lambda^+ - \mu^+, \rho \rangle - 1.$$

This characterization is a bit complicated, as we need to determine λ^+, μ^+ and the lengths of w^λ and w^μ . In the following two propositions, we give a simpler characterization of the cover relation $R_\alpha(\lambda) \xrightarrow{\alpha} \lambda$ when λ is α -regular.

Proposition 4.9. *Let μ, λ be two coweights in \check{P} such that $\mu = R_\alpha(\lambda)$ for some positive root α . Assume that $\langle \lambda, \alpha \rangle > 0$. Then $\mu \xrightarrow{\alpha} \lambda$ if and only if $w^\lambda \xrightarrow{\alpha} s_\alpha w^\lambda$.*

Proof. Observe that $w^\lambda < s_\alpha w^\lambda$ and $(w^\lambda)^{-1}(\alpha) \in \Phi^+$, which follow from the assumption that $\langle \lambda, \alpha \rangle > 0$. We also notice that $\lambda^+ = \mu^+$ in this case.

We first prove the direction “ \Leftarrow ”. It is enough to show that $w^\mu = s_\alpha w^\lambda$. By the formula (12), for any w and $\lambda \in \mathfrak{C}_w$, we always have

$$\ell(\tau_{-\lambda} w) \geq 2\langle \lambda^+, \rho \rangle - \ell(w).$$

In fact, the equality holds if and only if $w = w^\lambda$. We now show that $s_\mu = s_\alpha w^\lambda$ by contradiction. Assume that $s_\alpha w^\lambda \neq w^\mu$, implying the following inequality:

$$(31) \quad \ell(\tau_{-s_\alpha(\lambda)} s_\alpha w^\lambda) > 2\langle \lambda^+, \rho \rangle - \ell(s_\alpha w^\lambda) = 2\langle \lambda^+, \rho \rangle - \ell(w^\lambda) - 1.$$

On the other hand, we view $\tau_{-\lambda} w^\lambda$ as an element in the extended affine Weyl group \tilde{W}_{aff} . By the formula (20),

$$(\tau_{-\lambda} w^\lambda)^{-1}(\alpha) = (w^\lambda)^{-1}(\alpha) - \langle \lambda, \alpha \rangle \check{\delta}$$

is a negative affine root. It follows that $s_\alpha \tau_{-\lambda} w^\lambda < \tau_{-\lambda} w^\lambda$. In particular,

$$\ell(s_\alpha \tau_{-\lambda} w^\lambda) < \ell(\tau_{-\lambda} w^\lambda) = 2\langle \lambda^+, \rho \rangle - \ell(w^\lambda).$$

It contradicts with the inequality (31), since $s_\alpha \tau_{-\lambda} w^\lambda = \tau_{-s_\alpha(\lambda)} s_\alpha w^\lambda$.

The direction “ \Rightarrow ” is obvious. □

Proposition 4.10. *For any two coweights μ, λ in \check{P} with $\mu = R_\alpha(\lambda)$ for some positive root α , assume that $\langle \lambda, \alpha \rangle < 0$ and λ is α -regular. Then $\mu \xrightarrow{\alpha} \lambda$ if and only if $\ell(w^\lambda) - \ell(w^\mu) = -2\langle \check{\alpha}, w_\lambda(\rho) \rangle - 1$.*

Proof. By Lemma 4.7, $\mu = R_\alpha(\lambda) \in \mathfrak{C}_{s_\alpha w^\lambda}$. It follows that $\mu^+ = (w^\lambda)^{-1}(\lambda + \check{\alpha})$. Then

$$\langle \lambda^+ - \mu^+, 2\rho \rangle = -2\langle (w^\lambda)^{-1}(\check{\alpha}), \rho \rangle = -2\langle \check{\alpha}, w^\lambda(\rho) \rangle.$$

Hence $\mu \xrightarrow{\alpha} \lambda$ if and only if $\ell(w^\lambda) - \ell(w^\mu) = -2\langle \check{\alpha}, w_\lambda(\rho) \rangle - 1$. □

Note that if λ is $(r + 1)$ -regular, then $w^\mu = s_\alpha w^\lambda$; by the same argument as in Lemma 4.7, it is easy to check that if λ is $(r + 1)$ -regular, then $\mu = R_\alpha(\lambda) \in \mathfrak{C}_{s_\alpha w^\lambda}$ is 1-regular. Hence $w^\mu = s_\alpha w^\lambda$. This fact, along with Proposition 4.10, imply the following corollary.


Remark 4.11. *Prop.4.9 and Prop.4.10 are special cases of (LS, Proposition 4.1) when λ is super-regular in the sense of (LS). But the regularity for λ is much weaker in our propositions.*

Corollary 4.12. *With the same setup as in Proposition 4.10, assume that $\langle \lambda, \alpha \rangle < 0$ and λ is $(r + 1)$ -regular. If $R_\alpha(\lambda) \xrightarrow{\alpha} \lambda$, then $s_\alpha w^\lambda \xrightarrow{\alpha} w^\lambda$ if and only if $-(w^\lambda)^{-1}(\alpha)$ is a simple root.*

Now for any $\lambda \in \check{P}$, let $\Psi(\lambda)_\theta$ denote the subset of $\Psi(\lambda)$ consisting of μ such that $\mu \xrightarrow{\alpha} \lambda$ for some α . Geometrically, if $\mu \in \Psi(\lambda)_\theta$ then \bar{X}_μ is an irreducible divisor in \bar{X}_λ . In general, the set $\Psi(\lambda)_\theta$ consist of two types of elements $R_\alpha(\lambda)$, those for which $\langle \lambda, \alpha \rangle > 0$ and those for which $\langle \lambda, \alpha \rangle < 0$.

Example 4.13. Let G be of type A_2 . Let α_1 and α_2 be the two simple roots.

- (1) Take $\lambda = \check{\alpha}_1 + \check{\alpha}_2$. Then $\Psi(\lambda)_\theta = \{\check{\alpha}_1, \check{\alpha}_2\}$.
- (2) Take $\lambda = -(\check{\alpha}_1 + \check{\alpha}_2)$. Then $\Psi(\lambda)_\theta = \{0\}$.
- (3) Take $\lambda = -2(\check{\alpha}_1 + \check{\alpha}_2)$. Then $\Psi(\lambda)_\theta = \{-2\check{\alpha}_1 - \check{\alpha}_2, -\check{\alpha}_1 - 2\check{\alpha}_2, \check{\alpha}_1 + \check{\alpha}_2\}$.
- (4) Take $\lambda = -3\check{\alpha}_1$. Then $\Psi(\lambda)_\theta = \{2\check{\alpha}_2 - \check{\alpha}_1, -3(\check{\alpha}_1 + \check{\alpha}_2)\}$. The dimension of X_λ is 10. The following picture illustrates for example (4) the dimension of X_μ for all possible $\mu = R_\alpha(\lambda)$:



4.3. Braid relations on R -operators.

Proposition 4.14. *Let λ be a coweight in \check{P} , and let α, β be two positive roots in Φ .*

- (1) *If $\langle \alpha, \check{\beta} \rangle = 0$, then $R_\alpha R_\beta(\lambda) = R_\beta R_\alpha(\lambda)$.*

- (2) If $\langle \alpha, \check{\beta} \rangle = -1, \langle \beta, \check{\alpha} \rangle = -1$, and λ is away from the hyperplane $H_{\alpha+\beta,-1}$, then $R_\alpha R_\beta R_\alpha(\lambda) = R_\beta R_\alpha R_\beta(\lambda)$.
- (3) If $\langle \alpha, \check{\beta} \rangle = -1, \langle \beta, \check{\alpha} \rangle = -2$, and λ is away from the hyperplanes $H_{\alpha+\beta,-1}, H_{2\alpha+\beta,-1}$ and $H_{2\alpha+\beta,-2}$, then $R_\alpha R_\beta R_\alpha R_\beta(\lambda) = R_\beta R_\alpha R_\beta R_\alpha(\lambda)$.

Proof. We first show part (1). Assume that $\langle \alpha, \check{\beta} \rangle = 0$. We compare $R_\alpha R_\beta(\lambda)$ with $R_\beta R_\alpha(\lambda)$. There are four cases. If $\langle \lambda, \alpha \rangle \geq 0$ and $\langle \lambda, \beta \rangle \geq 0$, then the R operators coincide with the reflections s_α, s_β which commute, so we obviously have equality. Assume $\langle \lambda, \alpha \rangle < 0, \langle \lambda, \beta \rangle \geq 0$. Now we compute

$$R_\alpha R_\beta(\lambda) = R_\alpha s_\beta(\lambda) = s_\alpha s_\beta(\lambda) - \check{\alpha}.$$

The other expression evaluates to $s_\beta s_\alpha(\lambda) - s_\beta(\check{\alpha})$ and the two expressions are equal since $s_\beta(\check{\alpha}) = \check{\alpha}$. The case where $\langle \lambda, \alpha \rangle \geq 0$ but $\langle \lambda, \beta \rangle < 0$ is identical. Lastly if $\langle \lambda, \alpha \rangle < 0$ and $\langle \lambda, \beta \rangle < 0$ we get expressions $s_\alpha s_\beta(\lambda) - \check{\alpha} - \check{\beta}$ and $s_\beta s_\alpha(\lambda) - \check{\beta} - \check{\alpha}$, which again are equal.

The proof of part (2) and part (3) essentially follows by brute force computation. We include tables with all of the relevant information, as well as some sample computations.

In the tables below we make the following convention: if a square is blank, then the braid relation is satisfied. Only when the braid relation is not satisfied do we put anything in the second column. This is simply for readability.

Type A_2 : $\langle \alpha, \check{\beta} \rangle = -1, \langle \beta, \check{\alpha} \rangle = -1$

| Conditions | $R_\alpha R_\beta R_\alpha$ | $R_\beta R_\alpha R_\beta$ |
|---|--|--|
| $\langle \lambda, \alpha \rangle \leq -1, \langle \lambda, \beta \rangle \leq -1$ | $w(\lambda) - 2(\check{\alpha} + \check{\beta})$ | |
| $\langle \lambda, \alpha \rangle = -1, \langle \lambda, \beta \rangle = 0$ | $w(\lambda) - \check{\alpha} - \check{\beta}$ | |
| $\langle \lambda, \alpha \rangle = 0, \langle \lambda, \beta \rangle = -1$ | $w(\lambda) - \check{\alpha} - \check{\beta}$ | |
| $\langle \lambda, \alpha \rangle \geq 0, \langle \lambda, \alpha + \beta \rangle < -1$ | $w(\lambda) - 2\check{\alpha} - \check{\beta}$ | |
| $\langle \lambda, \alpha \rangle \geq 1, \langle \lambda, \alpha + \beta \rangle = -1$ | $w(\lambda) - 2\check{\alpha} - \check{\beta}$ | $w(\lambda) - \check{\alpha}$ |
| $\langle \lambda, \alpha + \beta \rangle \geq 0, \langle \lambda, \beta \rangle \leq -1$ | $w(\lambda) - \check{\alpha}$ | |
| $\langle \lambda, \alpha \rangle \geq 0, \langle \lambda, \beta \rangle \geq 0$ | $w(\lambda)$ | |
| $\langle \lambda, \beta \rangle \geq 0, \langle \lambda, \alpha + \beta \rangle < -1$ | $w(\lambda) - 2\check{\beta} - \check{\alpha}$ | |
| $\langle \lambda, \alpha + \beta \rangle = -1, \langle \lambda, \beta \rangle \geq 1$ | $w(\lambda) - \check{\beta}$ | $w(\lambda) - 2\check{\beta} - \check{\alpha}$ |
| $\langle \lambda, \alpha + \beta \rangle \geq 0, \langle \lambda, \alpha \rangle \leq -1$ | $w(\lambda) - \check{\beta}$ | |

Type B_2 : $\langle \alpha, \check{\beta} \rangle = -1, \langle \beta, \check{\alpha} \rangle = -2$

| Conditions | $R_\alpha R_\beta R_\alpha R_\beta$ | $R_\beta R_\alpha R_\beta R_\alpha$ |
|---|---|---|
| $\langle \lambda, \alpha \rangle \leq -1, \langle \lambda, \beta \rangle < 0$ | $w(\lambda) - 3\check{\alpha} - 4\check{\beta}$ | |
| $\langle \lambda, \alpha \rangle = 0, \langle \lambda, \beta \rangle = -2$ | $w(\lambda) - 2\check{\alpha} - 3\check{\beta}$ | |
| $\langle \lambda, \beta \rangle = 0, \langle \lambda, \alpha \rangle = -1$ | $w(\lambda) - 2\check{\alpha} - 2\check{\beta}$ | |
| $\langle \lambda, \beta \rangle = -1, \langle \lambda, \alpha \rangle = 0$ | $w(\lambda) - \check{\alpha} - 2\check{\beta}$ | |
| $\langle \lambda, \alpha + \beta \rangle < -1, \langle \lambda, \beta \rangle \geq 0$ | $w(\lambda) - 3\check{\alpha} - 3\check{\beta}$ | |
| $\langle \alpha + \beta \rangle = -1, \langle \lambda, \beta \rangle > 0$ | $w(\lambda) - 3\check{\alpha} - 3\check{\beta}$ | $w(\lambda) - 2\check{\alpha} - \check{\beta}$ |
| $\langle \lambda, \beta \rangle = 1, \langle \lambda, \alpha \rangle = -2$ | $w(\lambda) - 3\check{\alpha} - 3\check{\beta}$ | $w(\lambda) - 2\check{\alpha} - 2\check{\beta}$ |
| $\langle \lambda, \alpha + \beta \rangle \geq 0, \langle \lambda, 2\alpha + \beta \rangle < -2$ | $w(\lambda) - 2\check{\alpha} - 2\check{\beta}$ | |
| $\langle \lambda, \alpha \rangle \leq -2, \langle \lambda, 2\alpha + \beta \rangle = -2 \text{ or } -1$ | $w(\lambda) - 2\check{\alpha} - \check{\beta}$ | $w(\lambda) - \check{\alpha}$ |
| $\langle \lambda, \beta \rangle = 1, \langle \lambda, \alpha \rangle = -1$ | $w(\lambda) - \check{\alpha} - \check{\beta}$ | $w(\lambda) - \check{\alpha} - \check{\beta}$ |
| $\langle \lambda, 2\alpha + \beta \rangle \geq 0, \langle \lambda, \alpha \rangle \leq -1$ | $w(\lambda) - \check{\alpha}$ | |
| $\langle \lambda, \alpha \rangle \geq 0, \langle \lambda, \beta \rangle \geq 0$ | $w(\lambda)$ | |
| $\langle \lambda, \alpha \rangle \geq 0, \langle \lambda, 2\alpha + \beta \rangle < -2$ | $w(\lambda) - 2\check{\alpha} - 4\check{\beta}$ | |
| $\langle \lambda, \alpha \rangle > 0, \langle \lambda, 2\alpha + \beta \rangle = -2 \text{ or } -1$ | $w(\lambda) - \check{\alpha} - 3\check{\beta}$ | $w(\lambda) - 2\check{\alpha} - 4\check{\beta}$ |
| $\langle \lambda, 2\alpha + \beta \rangle \geq 0, \langle \lambda, \alpha + \beta \rangle < -1$ | $w(\lambda) - \check{\alpha} - 3\check{\beta}$ | |
| $\langle \lambda, \alpha + \beta \rangle = -1, \langle \lambda, 2\alpha + \beta \rangle \geq 2$ | $w(\lambda) - \check{\beta}$ | $w(\lambda) - \check{\alpha} - 3\check{\beta}$ |
| $\langle \lambda, \alpha + \beta \rangle = -1, \langle \lambda, 2\alpha + \beta \rangle = 0$ | $w(\lambda) - \check{\beta}$ | $w(\lambda) - \check{\alpha} - 2\check{\beta}$ |
| $\langle \lambda, \alpha + \beta \rangle \geq 0, \langle \lambda, \beta \rangle < 0$ | $w(\lambda) - \check{\beta}$ | |

As an illustration, we do an example calculation in type A_2 . Assume $\langle \lambda, \alpha \rangle \geq 0$ and $\langle \lambda, \alpha + \beta \rangle < -1$. Then we compute $R_\alpha R_\beta R_\alpha(\lambda)$. As a first step, we compute $\langle \lambda, \alpha \rangle$. By assumption $\langle \lambda, \alpha \rangle \geq 0$ so $R_\alpha(\lambda) = s_\alpha(\lambda)$. Thus

$$R_\alpha R_\beta R_\alpha(\lambda) = R_\alpha R_\beta(s_\alpha(\lambda)).$$

Next we compute $\langle s_\alpha(\lambda), \beta \rangle = \langle \lambda, s_\alpha(\beta) \rangle = \langle \lambda, \alpha + \beta \rangle$. By assumption, this is less than or equal to -1 . Thus

$$R_\alpha R_\beta(s_\alpha(\lambda)) = R_\alpha(s_\beta(s_\alpha(\lambda)) - \check{\beta}).$$

Lastly we compute

$$\langle s_\beta(s_\alpha(\lambda)) - \check{\beta}, \alpha \rangle = \langle s_\alpha(\lambda) + \check{\beta}, s_\beta(\alpha) \rangle = \langle \lambda + s_\alpha(\check{\beta}), s_\beta(\alpha + \beta) \rangle = \langle \lambda + \check{\alpha} + \check{\beta}, \beta \rangle.$$

The two assumptions at the beginning on λ force $\langle \lambda, \beta \rangle < -1$ so that $\langle \lambda + \check{\alpha} + \check{\beta}, \beta \rangle \leq -2 + 1 < 0$, so finally we see that $R_\alpha R_\beta R_\alpha(\lambda) = s_\alpha(s_\beta(s_\alpha(\lambda)) - \check{\beta}) - \check{\alpha}$. Simplifying this expression yields $R_\alpha R_\beta R_\alpha(\lambda) = s_{\alpha+\beta}(\lambda) - 2\check{\alpha} - \check{\beta}$ as desired. Computing $R_\beta R_\alpha R_\beta$ proceeds in a similar fashion.

We do another example in type B_2 where the braid relation holds. Let $\langle \lambda, 2\alpha + \beta \rangle \geq 0$ and $\langle \lambda, \alpha + \beta \rangle < -1$. We first compute $R_\alpha R_\beta R_\alpha R_\beta(\lambda)$. We first calculate $\langle \lambda, \beta \rangle$ which by assumption is < 0 . Thus the first simplification is

$$R_\alpha R_\beta R_\alpha R_\beta(\lambda) = R_\alpha R_\beta R_\alpha(s_\beta(\lambda) - \check{\beta}).$$

Next we compute $\langle s_\beta(\lambda) - \check{\beta}, \alpha \rangle = \langle \lambda + \check{\beta}, \beta + \alpha \rangle < 0$. So the next simplification is

$$R_\alpha R_\beta R_\alpha R_\beta(\lambda) = R_\alpha R_\beta((s_\alpha(s_\beta \lambda) - \check{\beta}) - \check{\alpha}).$$

The next pairing $\langle s_\alpha(s_\beta\lambda) - \check{\beta} - \check{\alpha}, \beta \rangle > 0$, so the next step is

$$R_\alpha R_\beta R_\alpha R_\beta(\lambda) = R_\alpha s_\beta(s_\alpha(s_\beta\lambda) - \check{\beta}) - \check{\alpha}.$$

One can then calculate the last pairing, see that it is ≥ 0 , and obtain finally that

$$R_\alpha R_\beta R_\alpha R_\beta(\lambda) = s_\alpha s_\beta(s_\alpha((s_\beta\lambda) - \check{\beta}) - \check{\alpha}) = s_\alpha s_\beta s_\alpha s_\beta(\lambda) - \check{\alpha} - 3\check{\beta}.$$

Computing the other side, $R_\beta R_\alpha R_\beta R_\alpha(\lambda)$ is much the same; we mention that the first two pairings will be positive and the last two negative, leading to

$$R_\beta R_\alpha R_\beta R_\alpha(\lambda) = s_\beta(s_\alpha(s_\beta s_\alpha(\lambda)) - \check{\alpha}) - \check{\beta} = s_\beta s_\alpha s_\beta s_\alpha(\lambda) - \check{\alpha} - 3\check{\beta}.$$

Thus, in this case the braid relation holds. \square

Remark 4.15. A summary of the data from the tables in the above proof is as follows: the “braid relations” for the R -operators hold in types A_2 , B_2 everywhere except at a certain set of critical lines. In the following w_0 denotes the longest element in Weyl group of each type. In type A_2 , the braid relations fail precisely when $\langle \lambda, \alpha + \beta \rangle = -1$ and $\lambda \notin \mathfrak{C}_{w_0}$. Similarly in B_2 , the braid relations hold except when $\langle \lambda, \alpha + \beta \rangle = -1$ and $\lambda \notin \mathfrak{C}_{w_0}$, or when $\langle \lambda, \beta + 2\alpha \rangle = -1$ or -2 and $\lambda \notin \mathfrak{C}_{w_0}$. Though we have not done all the calculations for G_2 , it seems to follow the same pattern: in particular the braid relations appear to hold in all cases, except the following: $\langle \lambda, \alpha + 2\beta \rangle = -1$, $\lambda \notin \mathfrak{C}_{w_0}$, or $\langle \lambda, 3\beta + 2\alpha \rangle = -1$ and $\lambda \notin \mathfrak{C}_{w_0}$.

Example 4.16. We do an example calculation in type B_2 where the braid relation fails. Let $\langle \lambda, \alpha + \beta \rangle = -1$ and let $\langle \lambda, 2\alpha + \beta \rangle \geq 2$. First we compute $R_\alpha R_\beta R_\alpha R_\beta$. As a first step we compute $\langle \lambda, \beta \rangle < 0$, so

$$R_\alpha R_\beta R_\alpha R_\beta(\lambda) = R_\alpha R_\beta R_\alpha(s_\beta(\lambda) - \check{\beta}).$$

Next we compute $\langle s_\beta(\lambda) - \check{\beta}, \alpha \rangle = \langle \lambda + \check{\beta}, \alpha + \beta \rangle$, which by assumption is 0, so we have

$$R_\alpha R_\beta R_\alpha R_\beta(\lambda) = R_\alpha R_\beta s_\alpha((s_\beta(\lambda) - \check{\beta})) = R_\alpha R_\beta(s_\alpha s_\beta(\lambda) - \check{\alpha} - \check{\beta}).$$

Computing

$$\langle s_\alpha s_\beta(\lambda) - \check{\alpha} - \check{\beta}, \beta \rangle = \langle s_\beta(\lambda) - \check{\beta}, \beta + 2\alpha \rangle = \langle \lambda + \check{\beta}, \beta + 2\alpha \rangle$$

which is forced to be ≥ 0 by our assumptions, we see that the next step is $R_\alpha(s_\beta s_\alpha s_\beta(\lambda) - \check{\alpha} - \check{\beta})$. The last computation is $\langle s_\beta s_\alpha s_\beta(\lambda) - \check{\alpha} - \check{\beta}, \alpha \rangle$; our assumptions force this to be positive, so our final result is

$$R_\alpha R_\beta R_\alpha R_\beta(\lambda) = s_\alpha s_\beta s_\alpha s_\beta(\lambda) - \check{\beta}$$

as desired. On the other hand, we compute $R_\beta R_\alpha R_\beta R_\alpha(\lambda)$. Since $\langle \lambda, \alpha \rangle > 0$ we must have $R_\beta R_\alpha R_\beta R_\alpha(\lambda) = R_\beta R_\alpha R_\beta s_\alpha(\lambda)$. Next we see that $\langle s_\alpha(\lambda), \beta \rangle > 0$, so we simplify further

$$R_\beta R_\alpha R_\beta R_\alpha(\lambda) = R_\beta R_\alpha(s_\beta s_\alpha(\lambda)).$$

Our assumptions force $\langle s_\beta s_\alpha(\lambda), \alpha \rangle < 0$, and so the next simplification is

$$R_\beta R_\alpha R_\beta R_\alpha(\lambda) = R_\beta(s_\alpha s_\beta s_\alpha(\lambda) - \check{\alpha}).$$

Lastly we compute $\langle s_\alpha s_\beta s_\alpha(\lambda) - \check{\alpha}, \beta \rangle < 0$. So we finally see

$$R_\beta R_\alpha R_\beta R_\alpha(\lambda) = s_\beta(s_\alpha s_\beta s_\alpha(\lambda) - \check{\alpha}) - \check{\beta} = s_\beta s_\alpha s_\beta s_\alpha(\lambda) - \check{\alpha} - 3\check{\beta}.$$

4.4. **Discussions on the moment polytope of \bar{X}_λ .** For any $\lambda \in \check{P}$, we define the moment polytope $\text{MP}(\lambda)$ of \bar{X}_λ as the convex hull of $\Psi(\lambda)$ in $\check{P} \otimes_{\mathbb{Z}} \mathbb{R}$. A part of our original motivation for this work was to explicitly understand the moment polytope $\text{MP}(\lambda)$, and understand whether all integral points (relative to λ) in $\text{MP}(\lambda)$ appear in $\Psi(\lambda)$.


Due to Theorem 4.1, if $\lambda \in \mathfrak{C}_w$ for some $w \in W$, then the part of the moment polytope inside the chamber \mathfrak{C}_w is easy to describe. In general, to describe a polytope, it suffices to describe the vertices of this polytope. For the polytope $\text{MP}(\lambda)$, all vertices are some special points in $\Psi(\lambda)$. Fix a chamber \mathfrak{C}_y . Let $\mathbb{M}_y(\lambda)$ denote the set of all coweights in $\Psi(\lambda)$ that are maximal in the chamber \mathfrak{C}_y . Using Theorem 4.1 again, we see that in any given chamber \mathfrak{C}_y the coweights of $\Psi(\lambda)$ appearing in chamber \mathfrak{C}_y are precisely those μ such that $\lambda' - \mu$ is a sum of coroots in $y(\Phi^+)$, for some $\lambda' \in \mathbb{M}_y(\lambda)$. Therefore the points in $\mathbb{M}_y(\lambda)$ are exactly the vertices of $\text{MP}(\lambda)$ in the chamber \mathfrak{C}_y .

We may use Proposition 4.8 to obtain candidates of points in $\mathbb{M}_y(\lambda)$. All of these points are obtained via $R_{\beta_1} \dots R_{\beta_k}(\lambda)$ so that $\mathfrak{C}_y = s_{\beta_1} \dots s_{\beta_k}(\mathfrak{C}_w)$, in other words $y = s_{\beta_1} \dots s_{\beta_k}w$. However in general it seems very difficult to determine which path of R -operators will provide the maximal vectors.

In the picture of the A_2 example below, we see five "paths" given by

$$R_\alpha R_\beta R_\alpha(\lambda), R_\beta R_\alpha R_\beta(\lambda), R_\alpha R_{\alpha+\beta} R_\beta(\lambda), R_\beta R_{\alpha+\beta} R_\alpha(\lambda), R_{\alpha+\beta}(\lambda),$$

where $\lambda = -3(\check{\alpha} + \check{\beta})$. Note that the last path $R_{\alpha+\beta}(\lambda)$ yields the maximal coweight $2(\check{\alpha} + \check{\beta})$, and all other candidates $\check{\alpha} + \check{\beta}$, $2\check{\beta} + \check{\alpha}$ and $2\check{\alpha} + \check{\beta}$ are in $\Psi(2\check{\alpha} + 2\check{\beta})$. Thus far this pattern has held in all our rank 2 experiments; in any given chamber \mathfrak{C}_w we have only ever observed one maximal coroot μ such that all other candidates $\mu' = R_{\beta_1} \dots R_{\beta_k}(\lambda) \in \Psi(\mu)$. In A_2 it seems to be the case (after much tedious calculation) that $R_\alpha R_\beta R_\alpha(\lambda) \in \Psi(R_{s_\alpha(\beta)})$, so in A_2 , to find the maximal candidates, one should replace $R_\alpha R_\beta R_\alpha$ with $R_{\alpha+\beta}$. On the other hand in B_2 there is an example where $R_\alpha(\lambda) \in \Psi(R_{\alpha+\beta} R_\beta R_{2\alpha+\beta}(\lambda))$, (note here $s_\alpha = s_{\alpha+\beta} s_\beta s_{2\alpha+\beta}$) so shorter expressions do not always do better.



One further remark is that in the above example, every integral coweight inside $\text{MP}(\lambda)$ is indeed contained in $\Psi(\lambda)$, where $\lambda = -3(\check{\alpha} + \check{\beta})$. If it were true in general that for any λ and for any $y \in W$, $\mathbb{M}_y(\lambda)$ contains at most one element (in other words, if each chamber has a unique "maximal" candidate if it exists), then this would imply the following

property: for every integral point μ inside $\text{MP}(\lambda)$, $\mu \in \Psi(\lambda)$. This property holds for $A_1 \times A_1$ and A_2 by naive calculation, and has held in every other example we have tried for B_2 . We would be interested in a proof or disproof of this property for general root systems. When λ is dominant, $\overline{X}_\lambda = \overline{\text{Gr}}_\lambda$. In this case, it is well-known that $\text{MP}(\lambda)$ is the convex hull of $\{w(\lambda) \mid w \in W\}$, and for every weight $\mu \in \text{MP}(\lambda)$ such that $\lambda - \mu \in \check{Q}$, $\mu \in \Psi(\lambda)$. The study of the moment polytopes of certain subvarieties in affine Grassmannian very often leads to interesting applications in representation theory, see (An, Kam).

REFERENCES

- [An] J. Anderson, A polytope calculus for semisimple groups. *Duke Math. J.* 116 (2003), no. 3, 567-588.
- [BB] A. Björner and F. Brenti, *Combinatorics of Coxeter groups*. Graduate Texts in Mathematics, 231. Springer, New York, 2005.
- [BD] A. Beilinson and V. Drinfeld, Quantization of Hitchin's integrable system and Hecke eigensheaves, available at www.math.uchicago.edu/mitya/langlands.
- [BGG] J. Bernstein, I. M. Gelfand, S. I. Gelfand. Schubert cells and cohomologies of spaces G/P . *Uspekhi Mat. Nauk* 38, No.3, 3-26(1973).
- [Bou] N. Bourbaki. *Lie Algebras and Lie Groups*, Chapters 4-6. Springer, Berlin, 2002.
- [BT] D. Bernard and J. Thierry-Mieg. Level one representations of the simple affine Kac-Moody algebras in their homogeneous gradations. *Comm. Math. Phys.* 111 (1987), no. 2, 181-246.
- [Fr] I. Frenkel, Two constructions of affine Lie algebra representations and boson-fermion correspondence in quantum field theory. *J. Funct. Anal.* 44 (1981), no. 3, 259-327.
- [FK] I. Frenkel and V. Kac. Basic representations of affine Lie algebras and dual resonance models. *Invent. Math.* 62 (1980/81), no. 1, 23-66.
- [FLM] I. Frenkel, J. Lepowsky and A. Meurman. *Vertex operator algebras and the Monster*. Pure and Applied Mathematics, 134. Academic Press, Inc., Boston, MA, 1988.
- [Hu1] J. Humphreys, *Introduction to Lie Algebras and Representation Theory*. Third Printing, Revised. Springer-Verlag 1980.
- [Hu2] J. Humphreys, *Reflection Groups and Coxeter Groups*. Cambridge studies in advanced mathematics volume 29. Cambridge University Press 1990.
- [IM] N. Iwahori and H. Matsumoto, On some Bruhat decomposition and the structure of the Hecke rings of p-adic Chevalley groups, *Inst. Hautes Études Sci. Publ. Math.* (1965), no. 25, 5-48.
- [Io1] B. Ion, Nonsymmetric Macdonald Polynomials and Demazure Characters. *Duke Math. J.* Vol. 116, No. 2 (2003), 299-318.
- [Io2] B. Ion. A weight multiplicity formula for Demazure modules. *Int. Math. Res. Not.* 2005, no. 5, 311-323.
- [Io3] B. Ion. Nonsymmetric Macdonald polynomials and matrix coefficients for unramified principal series. *Advances in Mathematics* 201 (2006), 36-62.
- [Io4] B. Ion. Standard bases for affine parabolic modules and nonsymmetric Macdonald polynomials.

- [Ka] V. Kac, Infinite-dimensional Lie algebras. Third edition. Cambridge University Press, Cambridge, 1990.
- [Kam] J. Kamnitzer, Mirković-Vilonen cycles and polytopes. *Ann. of Math. (2)* 171 (2010), no. 1, 245-294.
- [KT] M. Kashiwara and T. Tanisaki, Parabolic Kazhdan-Lusztig polynomials and Schubert varieties. *J. Algebra* 249 (2002), no. 2, 306-325.
- [Ku] S. Kumar, Kac-Moody Groups, their Flag Varieties and Representation Theory. *Progress in Mathematics*, 204. Birkhäuser Boston, Inc., Boston, MA, 2002.
- [LS] T. Lam and M. Shimozono, Quantum cohomology of G/P and homology of affine Grassmannian. *Acta Math.* 204 (2010), no. 1, 49-90.
- [Mac] I. G. Macdonald. *Affine Hecke Algebras and Orthogonal Polynomials*. Cambridge University Press, 2003.
- [MV] I. Mirković and K. Vilonen. Geometric Langlands duality and representations of algebraic groups over commutative rings. *Ann. of Math. (2)* 166 (2007), no. 1, 95-143.
- [Sch] C. Schwer. Galleries, Hall-Littlewood polynomials, and structure constants of the spherical Hecke algebra. *Int. Math. Res. Not.* 2006, Art. ID 75395, 31 pp.
- [St] J. Stembridge, The Partial Order of Dominant Weights. *Advances in Mathematics*, Volume 136, Issue 2, June 1998. Pages 340-364.
- [Stg] R. Steinberg. *Lectures on Chevalley groups*. University Lecture Series volume 66 (2010).
- [Zhu] X. Zhu, An introduction to affine Grassmannians and the geometric Satake equivalence. *Geometry of moduli spaces and representation theory*, 59-154, IAS/Park City Math. Ser., 24, Amer. Math. Soc., Providence, RI, 2017.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL, CHAPEL HILL, NC 27599-3250, U.S.A.

Email address: marmarc@live.unc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL, CHAPEL HILL, NC 27599-3250, U.S.A.

Email address: jiuzu@email.unc.edu